PLANT DISEASE REPORTER INDEX

Issued by

CROPS RESEARCH DIVISION

Plant Industry Station
Volume 50

Beltsville, Maryland
Number 13

AUTHOR INDEX

ADAMS, R. E., (545)
ADLERZ, W. C., (602)
AFANASIEV, M. M., 928
AGNIHOTRI, J. P., 921
AGRIOS, GEORGE N., 151
AICHELE, MURIT D., (3)
ALLAM, E. K., 761
ALLAN, R. E., (205)
ALLEN, ROSS M., 145
ALLEN, W. WAYNE, 622
ANDERSON, P. M., (363)
ARMSTRONG, G. M., 391
ARMSTRONG, JOANNE K., (391)
ARNY, D. C., (12), (377), 380
ATLEE, C. B., Jr., (611)
AULAKH, KIRPAL S., 314

BACHELDER, S., (294)
BAGGETT, J. R., 532
BAIN, DOUGLAS C., 770, (770)
BAJAJ, B. S., (299)
BALDRIDGE, JOE D., (753)
BALL, ELLEN M., (951)
BANTTARI, E. E., 17
BARBE, G. DOUGLAS, 768
BARD, G., (722)
BARKAI-GOLAN, RIVKA, 874
BARKER, K. R., (420), (954), 957
BARNES, D. K., (484)
BARNES, GEORGE L., 599
BASHAW, E. C., (622)
BATRA, S. K., 299
BAUER, A. H., (6)
BEAN, GEORGE A., 942
BEARDMORE, R. W., (798)
BEGA, ROBERT V., (181), (245), 439, 832
BELL, DURHAM K., 162
BERAHA, L., (283)
BERGER, RICHARD D., (701)
BERKENKAMP, B., 390
BERRY, F. H., 45
BIESBROCK, J. A., 360
BIRCHFIELD, WRAY, 149, (500)
BLODGETT, EARLE C., 3
BOND, WILLIAM P., 325
BOONE, DONALD M., (539), 543, (957)
BORDERS, HUEY I., 847
BOYCE, JOHN S., Jr., (360), 482
BOYLE, LYTTON W., (810)
BRAKKE, MYRON K., (951)
BRAUN, ALVIN J., 783, (877)
BRAVERMAN, SAMUEL W., 321, 488, 849
BRAVO, LUIS E., (101)
BRETZ, T. W., (45)
BRINKERHOFF, L. A., 323
BROWDER, L. E., 673, (756)
BROWN, G. ELDON, 860
BRZESKI, M. W., (584)
BUCHHOLTZ, W. F., 88
BUGBEE, W. M., 576
BUTLER, E. E., 796
BYRD, D. W., Jr., 954
BYRNE, T. G., (751)

CALVERT, OSCAR H., (753)
CAMPBELL, W. A., (393)
CANNON, T. F., (664)
CARLSON, L. W., 539
CARPENTER, JOHN B., (145)
CASTIBLANCO, LUIS E., (101)
CEPONIS, MICHAEL J., 222
CHAKRAVARTI, B. P., 701
CHAMBERS, ALBERT Y., (608)
CHI, C. C., 451, 695
CHILDERS, W. R., (695)
CHILDS, J. F. L., 81
CLARK, W. A., (502)
COHEN, ELIAHOU, (251), (867)
COHEN, Y., (635)
COHN, E., 711
COLE, H., Jr., 494
COLLINS, DON D., 45
COMBS, R. L., Jr., (570)
CONVERSE, R. H., 351
COOK, A. A., (761)
COOPER, R. L., (557)
COOPER, W. E., 136
CORDELL, CHARLES E., 589
COUNTER, BEN F., (371)
COVEY, R. A., (645)
COVEY, RONALD P., Jr., 787
COX, R. S., 124, (354), 699
COYNE, D. P., 29
CRANE, J. L., 464
CRITTENDEN, H. W., (464), 910
CROSIER, WILLARD F., (321)
CROSS, J. E., 557, (560)
CROSSAN, D. F., (84), (647)
CULP, T. W., 35

DAFT, GILBERT C., 493
DALE, J. L., 441
DAVIS, D. R., 906
DAVIS, J. R., (53)
DÍAZ, FRANCISCO, (422)
DICKENS, LESTER E., (371)

PLANT DISEASE REPORTER INDEX

Issued by

CROPS RESEARCH DIVISION

Plant Industry Station
Volume 50

Beltsville, Maryland
Number 13

AUTHOR INDEX

ADAMS, R. E., (545)
ADLERZ, W. C., (602)
AFANASIEV, M. M., 928
AGNIHOTRI, J. P., 921
AGRIOS, GEORGE N., 151
AICHELE, MURIT D., (3)
ALLAM, E. K., 761
ALLAN, R. E., (205)
ALLEN, ROSS M., 145
ALLEN, W. WAYNE, 622
ANDERSON, P. M., (363)
ARMSTRONG, G. M., 391
ARMSTRONG, JOANNE K., (391)
ARNY, D. C., (12), (377), 380
ATLEE, C. B., Jr., (611)
AULAKH, KIRPAL S., 314

BACHELDER, S., (294)
BAGGETT, J. R., 532
BAIN, DOUGLAS C., 770, (770)
BAJAJ, B. S., (299)
BALDRIDGE, JOE D., (753)
BALL, ELLEN M., (951)
BANTTARI, E. E., 17
BARBE, G. DOUGLAS, 768
BARD, G., (722)
BARKAI-GOLAN, RIVKA, 874
BARKER, K. R., (420), (954), 957
BARNES, D. K., (484)
BARNES, GEORGE L., 599
BASHAW, E. C., (622)
BATRA, S. K., 299
BAUER, A. H., (6)
BEAN, GEORGE A., 942
BEARDMORE, R. W., (798)
BEGA, ROBERT V., (181), (245), 439, 832
BELL, DURHAM K., 162
BERAHA, L., (283)
BERGER, RICHARD D., (701)
BERKENKAMP, B., 390
BERRY, F. H., 45
BIESBROCK, J. A., 360
BIRCHFIELD, WRAY, 149, (500)
BLODGETT, EARLE C., 3
BOND, WILLIAM P., 325
BOONE, DONALD M., (539), 543, (957)
BORDERS, HUEY I., 847
BOYCE, JOHN S., Jr., (360), 482
BOYLE, LYTTON W., (810)
BRAKKE, MYRON K., (951)
BRAUN, ALVIN J., 783, (877)
BRAVERMAN, SAMUEL W., 321, 488, 849
BRAVO, LUIS E., (101)
BRETZ, T. W., (45)
BRINKERHOFF, L. A., 323
BROWDER, L. E., 673, (756)
BROWN, G. ELDON, 860
BRZESKI, M. W., (584)
BUCHHOLTZ, W. F., 88
BUGBEE, W. M., 576
BUTLER, E. E., 796
BYRD, D. W., Jr., 954
BYRNE, T. G., (751)

CALVERT, OSCAR H., (753)
CAMPBELL, W. A., (393)
CANNON, T. F., (664)
CARLSON, L. W., 539
CARPENTER, JOHN B., (145)
CASTIBLANCO, LUIS E., (101)
CEPONIS, MICHAEL J., 222
CHAKRAVARTI, B. P., 701
CHAMBERS, ALBERT Y., (608)
CHI, C. C., 451, 695
CHILDERS, W. R., (695)
CHILDS, J. F. L., 81
CLARK, W. A., (502)
COHEN, ELIAHOU, (251), (867)
COHEN, Y., (635)
COHN, E., 711
COLE, H., Jr., 494
COLLINS, DON D., 45
COMBS, R. L., Jr., (570)
CONVERSE, R. H., 351
COOK, A. A., (761)
COOPER, R. L., (557)
COOPER, W. E., 136
CORDELL, CHARLES E., 589
COUNTER, BEN F., (371)
COVEY, R. A., (645)
COVEY, RONALD P., Jr., 787
COX, R. S., 124, (354), 699
COYNE, D. P., 29
CRANE, J. L., 464
CRITTENDEN, H. W., (464), 910
CROSIER, WILLARD F., (321)
CROSS, J. E., 557, (560)
CROSSAN, D. F., (84), (647)
CULP, T. W., 35

DAFT, GILBERT C., 493
DALE, J. L., 441
DAVIS, D. R., 906
DAVIS, J. R., (53)
DÍAZ, FRANCISCO, (422)
DICKENS, LESTER E., (371)

DICKERSON, O. J., 396
DICKSON, R. C., 305
DIMOCK, A. W., 751
DISHON, IRENE, (21)
DOBIE, N. D., 426
DOEPEL, R. F., 494
DOOLEY, H. L., (97)
DOOLING, O. J., 862
DOTTA, D., (439)
DOUGLAS, W. A., (570)
DUBIN, H. JESSE, 280
DUKES, P. D., (275), (444), (520), (836)
DUNLEAVY, JOHN, (598)
DWINELL, L. D., (406)

ECK, PAUL, (346)
ECKESS, E. D., (571)
EDGERTON, L. J., (240)
EDMUNDS, J. E., 478
EL-HAIDARI, H. S., 960
ELLINGBOE, A. H., (190)
ENGEL, R. E., (703)
ENGLISH, HARLEY, (53)
EPPS, JAMES M., 608
EPSTEIN, A. H., 167, 420
ERWIN, D. C., (930)
EVERETT, PAUL H., (340)

FARRELL, K. M., (478)
FATHI, SAAD M., (401)
FEDER, WILLIAM A., 429
FILER, T. H., Jr., 264, 525, 751
FISHER, FRAN. E., 357
FLEISCHMANN, G., 750
FLIEGEL, PETER, 240
FORD, RICHARD E., (88), (814)
FOSKETT, RICHARD L., (309)
FOSTER, R. E., (49), 79
FRAZIER, D. H., (510)
FRAZIER, N. W., 318
FRAZIER, W. A., (532)
FREDERIKSEN, R. A., 840
FREEMAN, T. E., 292
FREITAG, J. H., (318)
FREYRE, R. H., (484)
FRIDLUND, PAUL R., 740, 902
FROSHEISER, F. I., 435
FULTON, N. D., (350)
FULTON, ROBERT W., (613)
FUNK, C. R., (294), (703)
FUTRELL, MAURICE C., (373), 606, 641, 828

GARREN, R. E., (843)
GARRETT, WILEY N., (839)
GASKILL, JOHN O., 457
GASKIN, T. A., 430
GAUTREAU, E. J., (419)
GHAFFAR, ABDUL, 171
GILMER, R. M., 461
GOLDEN, A. MORGAN, (645), (798)
GOOCH, F. S., 525
GOODING, G. V., Jr., 308
GOTH, R. W., 33, 110
GRAVES, ALFRED A., Jr., (482)
GREATHEAD, A. S., 327
GRIERSON, W., (860)
GRIFFITHS, D. A., 116, 261
GROVES, A. B., (461)
GUDAUSKAS, ROBERT T., 885
GUENGERICH, H. W., 367
GUTTER, Y., 782

HAASIS, F. A., (664)
HAGEDORN, D. J., (91), (578)
HALISKY, P. M., 294, 703
HALL, D. H., 793, (796)
HALLIWELL, ROBERT S., 75
HAMBLEN, M. L., (15)
HAMILTON, R. I., (822)
HAMMONS, R. O., (520)
HANLIN, RICHARD T., 306
HANSEN, A. JUERGEN, 229
HARDISON, JOHN R., 624
HARLAN, D. P., 548
HARMON, DANIEL L., (502)
HARMON, S. A., (443)
HARRISON, A. L., 855
HARRISON, MONTY D., 897
HARROW, K. M., (730)
HARTWIG, E. E., (104)
HARVEY, GEORGE M., 554
HASAN, S., 764
HASTINGS, K. M., (910)
HELTON, A. W., (773)
HENDRIX, FLOYD F., Jr., 393
HENDRIX, J. WALTER, (369)
HENSON, LAWRENCE, (218)
HERNANDEZ, TEME P., (500)
HERR, L. J., 777
HIBBEN, C. R., 905
HILTY, J. W., 427
HIMELICK, E. B., (473)
HINDS, THOMAS E., 741
HIRATSUKA, Y., 419
HOBBS, C. D., 373
HOCKING, D., 593
HOES, J. A., 62, 154
HOLDEMAN, Q. L., (318)
HOPKINS, DONALD L., (218)
HOPPE, PAUL E., 377
HORNE, C. WENDELL, 869
HORNER, C. E., 97
HOWELL, R. W., (104)
HSI, DAVID C. H., 175, 352
HUBER, D. M., 852
HULLUKA, MENGISTU, 631
HUTCHINS, LEE M., 219
HUTCHINS, P. C., (429)
HYRE, R. A., 734

IIDA, T. T., (453)
IZADPANAH, K., 779

JEDLINSKI, H., (380)
JENKINS, L., (548)
JENKINS, S. F., Jr., (275), 520, 836
JENSEN, HAROLD J., 923
JENSEN, RAY E., 810
JOHNSON, D. E., (133), (611), 915
JOHNSON, E. M., 218
JOHNSON, LOWELL B., 108
JOHNSON, R. E., (81)
JOHNSTON, C. O., 756
JONES, JOHN PAUL, 340, 707
JORGENSON, E. C., 650
JOSEPHSON, L. M., (427)

KAHAN, R. S., (874)
KAIS, A. G., (388), 842
KALLIO, ARVO, 69
KALOOSTIAN, G. H., (270)
KANTZES, J. G., (351)
KAPPELMAN, A. J., Jr., 655
KATSANOS, R. A., 287
KATZIR, R., (594)
KEEN, NOEL T., (192)
KENNEDY, B. W., (557), 560
KEPLINGER, JOSEPH A., (783), (877)
KERNKAMP, M. F., (576), (800)
KHANNA, R. N., (127)
KILBY, W. W., (54), (717)
KILPATRICK, R. A., (622), 789
KLINE, D. M., (382)
KOENIGS, J. W., 528
KOMMEDAHL, THOR, 26
KOZELNICKY, GEORGE M., 839, 840
KRIKUN, J., (711)
KRULL, CHARLES F., 101
KUNTZ, JAMES E., (677), (736)

LABORDE, J. A., (947)
LACKEY, C. F., 479
LAIRD, E. F., Jr., (305)
LAMBE, R. C., (88)
LAMBERT, J. W., (557)
LANGDON, K. R., (822)
LATHAM, A. J., 469
LAURENT, T. H., 921
LAUT, JOHN G., 208
LAWRENCE, J. J., (208)
LAYNE, R. E. C., 112
LEAR, BERT, 132, 133, 611, 858, (915)
LEATH, K. T., 312
LEBEN, CURT, (493)
LEEPER, PAUL W., (460)
LEWIS, MARJORIE H., 681
LEWIS, W. A., (54), (717)
LIM, W. C., (116), (261)
LITTRELL, R. H., 882
LIVINGSTON, CLARK H., (897)
LOEGERING, WILLIAM Q., 502
LORIO, PETER L., Jr., 596
LUCHT, DONALD D., (126)
LUEPSCHEN, N. S., 565
LUKE, H. H., (292)

MacKENZIE, D. R., (6), 363, 814
MacSWAN, I. C., (267)
MAIER, CHARLES R., 301
MANNING, W. J., 84, 647
MARLATT, ROBERT B., 687, 689
MARTENS, J. W., 12
MARTIN, JEAN E., 822
MARTIN, NEIL E., 369
MARTIN, W. J., 500
MATHRE, D. E., 930
MATHUR, S. B., 619
MAXIE, E. C., (327)
McCARTER, STATES M., (885)
McCARTNEY, WILLIAM O., (768)
McCLAIN, L. A., 112
McCLELLAN, W. D., 725
McCOLLOCH, LACY P., 178, 348
McCRUM, R. C., 40
McDANIEL, M. C., 350
McDONALD, H. H., (947)
McELROY, F. D., 581
McINTOSH, D. L., 267
McKINNEY, H. H., 951
McLEAN, D. M., 871
McWHORTER, F. P., (532)
MEYER, R. W., (938)
MICHEL, L. J., 935
MILBRATH, J. A., 59
MILLECAN, ARTHUR A., 211
MILLER, D. R., (439)
MILLER, J. W., 187
MILLER, LAWRENCE I., 139
MILLER, P. M., 201, 506
MILLER, P. W., (426), 843
MILLER, PAUL R., 803
MILLIKAN, D. F., (367)
MINK, G. I., (461)
MISHRA, M. D., (619)
MITCHELL, J. E., 91
MIYAGAWA, S. T., (132), (133), (611)
MOHAMED, HOSNI A., 401
MOLLER, W. J., 53
MONCRIEF, JAMES B., (840)
MONEIM, AHMED T. ABDEL, (401)
MOORE, D. M., (910)
MOORE, J. N., 105
MORGAN, FREDERIC L., 104, 598
MORRIS, C. L., 510
MORTON, D. J., 96, 275, (443), 444, (836)
MOTSINGER, R. E., 686
MUELLER, WALTER C., 899
MUKHERJEE, A. K., 88

NATTI, J. J., 614
NEELY, DAN, 473

NELSON, MERRITT R., (629), 947
NELSON, R. R., 382
NETZER, DAVID, 21, 594
NEWHALL, ALLAN G., 422
NICHOLS, LESTER P., 515
NICKLE, W. R., 798
NIELSEN, L. W., 730
NITZANY, F. E., 158
NORTON, DON C., 523
NORTON, J. D., (661)
NUSBAUM, C. J., (954)

O'BANNON, JOHN H., (512)
OHMAN, JOHN H., 402
OLOFSSON, JAN, 257
ORELLANA, R. G., 7
ORLOB, GERT B., 819, 917
ORR, CALVIN C., (149), 645
OSHIMA, NAGAYOSHI, 309, 371
OSTAZESKI, STANLEY A., 825
OTTERBACHER, A. G., (65)
OU, S. H., (453)
OUELLETTE, G. B., 722

PAI, H. SURENDRANATH, 313, 518
PALMER, L. T., 234
PALMITER, D. H., (783), 877
PAPPELIS, A. J., (287)
PARACER, S. M., 584
PARKER, K. G., (240)
PARRIS, G. K., 243
PARTRIDGE, A. D., 497
PATEL, K. P., (26)
PATWARDHAN, P. G., 709
PAULUS, A. O., (930)
PAVGI, M. S., 119
PEACHER, P. H., (862)
PEEPLES, J. L., Jr., 770
PENDERY, W., (793)
PÉREZ, J. ENRIQUE, 569
PETERSON, GLENN W., (618), 659
PETERSON, ROGER S., (741), 744
PETTIT, ROBERT E., 753
PHELPS, WILLIAM R., 224, 736
PIERCE, WALTER H., (238)
PIERSON, CHARLES F., 885, 913
PINCKARD, J. A., 172, 254
PINE, T. S., 64, 289
PIRONE, THOMAS P., (325)
PITRE, H. N., 409, 570
PITTS, RALPH, 238
PLAUGHER, NOAH H., (22)
POWELL, J. M., 144
POWELSON, R. L., 806
POWERS, H. R., Jr., 432
PRASAD, KRISHNA, (661)
PRASAD, N., (385), (921)
PROVVIDENTI, R., (337)
PURDY, LAURENCE H., 205

QUIAOIT, AVELINA R., 613
QUICK, CLARENCE R., 550

RAABE, ROBERT D., 411, 519, 709
RAHMAN, M. ABDEL, (761)
RANGARAJAN, M., (701)
RASKI, D. J., (27)
RAVENSCROFT, A. V., (930)
RAYCHAUDHURI, S. P., (88)
REEVES, E. L., (787)
REID, C. P. P., 345
REYES, LUCAS, (840)
REYNOLDS, HAROLD W., (149), 512
RICH, A. E., (40)
RICH, SAUL, 721
RICO, GABRIEL, (101)
RIFFLE, JERRY W., 126, 677
RIGGS, R. D., 15
RIRIE, D., (327)
RIVERA, C. T., 453
ROBAYO, GABRIEL, (101)
ROBERTS, B. J., (332), 571, (631)
ROGERS, W. E., (569)
ROHDE, R. A., (489)
ROHRBACH, K. G., 773
ROMIG, R. W., (571)
RONCADORI, R. W., (432)
ROSENBERG, DAN Y., (768)
ROSENKRANZ, E. E., (409)
ROSS, ARNOLD, (736)
ROSS, ELDON W., 527
ROTEM, J., 635
ROTHMAN, PAUL G., 692
ROWELL, J. B., 800
RUPPEL, E. G., 484

SAINT-LAURENT, M., (356)
SAMBORSKI, D. J., (750)
SASSER, J. N., 664
SCHAFER, JOHN F., (108)
SCHEFFER, R. P., (190)
SCHENCK, N. C., 327, 602
SCHIEBER, EUGENIO, 694
SCHIFFMANN-NADEL, MINA, 251, 867
SCHNATHORST, W. C., 168
SCHNEIDER, BERNARD A., 934
SCHNEIDER, C. L., (457)
SCHNEIDER, HENRY, (270)
SCHOENEWEISS, DONALD F., 13, 196, 446
SCHREIBER, LAWRENCE R., 122, (573), 808
SCHROEDER, W. T., 337
SCHUSTER, M. L., (29)
SCIARONI, R. H., (133)
SECHLER, DALE T., (292), (906)
SEELY, C. I., (852)
SEHGAL, O. P., 862
SHANER, G. E., (806)
SHANKS, CARL H., Jr., 586
SHAUGHNESSY, LYLE, (29)
SHAW, C. G., (399)
SHAY, J. R., (461)
SHEPHERD, R. J., (793)

SHER, S. A., (581)
SILL, W. H., Jr., 11, (234)
SIMONS, M. D., (935)
SIMPSON, W. R., 215
SINCLAIR, J. B., (350), (681), (686)
SINCLAIR, W. A., 406
SINGH, R. D., 385
SINGH, R. S., 127, 668
SITARAMAIAH, K., (668)
SITTON, B. G., (54)
SKOOG, H. A., 627
SLEETH, BAILEY, 460
SMERLIS, E., 356
SMILEY, J. H., (218)
SMITH, C. H., (650)
SMITH, M. A., (283)
SMITH, N. A., 190
SMITH, RICHARD S., Jr., 181, 245, 249, (439), 832
SNOW, G. A., 388
SORENSEN, E. L., (731)
SOUSA, A. A., (424)
SOWELL, GROVER, Jr., 661
SPRINGER, J. K., (715)
SPURR, HARVEY W., Jr., 424
STALEY, JOHN M., (280)
STAMBAUGH, WILLIAM J., (589)
STANDIFER, LEON C., (172)
STAPLES, ROBERT, (951)
STEGALL, W. A., Jr., 476
STEWART, D. M., (312), 332
STIPES, R. JAY, 573, (808)
STOKES, D. E., 822
STONE, WILLIAM J. H., 414, 629
STRETCH, A. W., (346), 715
STUTEVILLE, D. L., 731
SUTHERLAND, J. R., 545
SUTTON, B. C., (208)

TARRY, JERRY C., (181), 399
TAYLOR, CARLTON F., (6), 701
THAMES, WALTER H., Jr., (869)
THOMAS, C. A., (35)
THOMPSON, D. L., (655)
THOMPSON, MARY RUTH, (65), (779)
THORNBERRY, H. H., 65, (779)
THYR, B. D., (96)
TIWARI, V. P., (619)
TODD, F. A., (308)
TOLER, R. W., 316
TOOLE, E. RICHARD, 160, 552, 945
TORIO, JOYCE C., 346
TOWERS, BARRY, 747
TRAYLOR, JACK A., (278)
TROLL, J., 489
TROUTMAN, JOSEPH L., (139)
TRUE, R. P., (545)
TSAO, PAMELA W., 270
TYSON, I. H., (62)

VAGO, C., (764)
VALBUENA, LUIS A., (101)
VALCARCE, ARLAND C., (446)
van der ZWET, T., 54, 717
Van GUNDY, S. D., (581)
VARNEY, E. H., (351)
von SCHMELING, B., 645

WAGNON, H. KEITH, 278
WATSON, ALICE J., (348)
WATSON, R. D., (469), (852)
WEBB, R. E., 49
WEBER, RAY, (224)
WEBSTER, ORRIN J., (606), (641), (828)
WEHLBURG, C., 354, 938
WEIDENSAUL, T. CRAIG, 22
WEINER, A., 27
WERNHAM, C. C., 6, (363), (814)
WESTER, R. E., (316)
WILLIAMS, HAROLD E., (278)
WILLIAMS, PAUL H., 192
WILLIAMS, W. T., (800)
WILSON, E. E., 182
WOLFENBARGER, D. O., 908
WORLEY, R. E., 443
WRIGHT, W. R., 283
WYSONG, DAVID S., 618

YANG, SHAW-MING, 578
YARWOOD, C. E., 639, 791

ZAIGER, D., 892
ZENTMYER, G. A., (892)
ZIMMERMAN, L. H., (35)
ZUCKERMAN, B. M., (584)

SUBJECT INDEX

Abies sp.: previously undescribed branch and stem canker with severe resinosis on balsam fir in Anticosti Island, 722
--- bracteata: 181
Acanthorhynchus vaccinii: 539
Acer saccharum: circumstantial evidence suggests that nematodes of several genera may contribute to incidence of maple blight and dieback in Wisconsin, 677
Aceria tulipae: 819; 917; 951
Adenostema fasciculatum: possible reservoir of a virus which resembles tobacco streak, recovered from dodder growing on this shrub in southern California, 479
Alabama: 885

Alaska: 69; 921
Albizia julibrissin: 945
Alfalfa: damping-off, effects of 4 fungicides on seedlings and on controlling Rhizoctonia solani in sand, 435; leaf spot and damping-off distribution in Kansas, 731; Phytophthora root rot found in Canada for first time, potential of disease and comparison of Canadian and American isolates, 451; residue as a soil amendment for control of onion bulb rotting, 469; survey of fungi associated with wilts and root rots and an assess of damage to alfalfa in Ontario, 695
Almond (see Prunus)
Alternaria brassicicola: 764
--- dauci: 594
--- ricini: 35
--- tenuis: causes black core rot of mandarin oranges, 127
Amendments: alfalfa residue, to reduce onion infection and bulb rotting, 469; soil, in India the addition of oilcakes of margosa, castor, and peanut considerably reduced root-knot nematode populations in soil cropped to okra and tomato, 669
Ampelopsis aconitifolia: new host for Xiphinema index, 27
Anguina spp.: 45
Announcements: 280; 624
Anthracnose: of Alpine currant, variation in Pseudopeziza ribis isolates from different hosts, 196
---: of cucumber, control, 340
---: of Dioscorea alata [winged yam], efficacy of different fungicides for control, 385
---: on Easter lily blossoms, caused by black-scale Colletotrichum, 751
---: new disease: of Barbados cherry in India, 518; of Phyllanthus acidus caused by Colletotrichum in India, 313
---: spotted, of hybrid tea rose petals, possibly first report of Sphaceloma on petals of Rosa, 934
---: stem, of lima bean in the South, 1965, 316
Antibiotics: of 4 tested in combination with dimethyl sulfoxide, none gave satisfactory control of Dutch elm disease, 573; effects of several on oak wilt organism in the field, 736
---: agricultural streptomycin (100 ppm) useful spray for tomato bacterial spot disease in south Florida, 699
---: cycloheximide and phytoactin: not effective in controlling blister rust cankers in eastern white pine, 224; evaluated for control of western gall rust and aspen cankers on lodgepole and ponderosa pines, 741
---: Pentaene G8 completely inhibits seedborne fungi of chili and reduces stem rust pustules on wheat, 299
---: semicarbazone and methyl hydrazone derivatives of cycloheximide evaluated for control of comandra rust on lodgepole pine, 528
Aphanomyces euteiches: 91
Aphelenchoides besseyi: 689
Aphis gossypii: 305
Apple: growth of newly transplanted trees affected in many ways by chemical soil treatments and plastic mulches used at planting time, 201; lenticel injury resulted on Golden Delicious apples stored in molded paper pulp trays and on fruit covered with filter paper both containing urea-formaldehyde wet-strength resins, but not in those containing polyamide resins, 885
---: canker: a canker disease of red delicious in Guatemala resembles blister bark disease, 694; crater canker, bark disease of Golden Delicious trees, cause unknown, extent of decline is proportional to amount of trunk injury, 3; European canker in California, importance of ascospores and macroconidia, susceptibility of leaf scars, and of rain, 182
---: nematodes, survey of orchards in Hudson Valley, 1956-1965, 783
---: powdery mildew, control by 2 new sulfites, 645
---: rots: blue-mold rot, fungicides for control, 913; collar rot of 1 to 7 year old trees propagated on East Malling II and Malling Merton 104 rootstocks, all growing in one block, 267; fruit rot caused by Phialophora malorum, first report in Pennsylvania, 701; perennial canker rot of, in West Virginia and Pennsylvania, 348
---: severe non-parasitic russeting and dimpling possibly result of frost injury or low temperatures during early fruit development, 151
---: stem-cavity browning associated with brown core of stored McIntosh, and possibly both disorders have common cause, 178
---: viruses: transmission studies with selective Malus indicators, 40; virus indexing procedures to be followed by the IR-2 Pome Fruit

Virus Subproject, under auspices of American Phytopathological Society, 461
Apricot (see also Prunus): bacterial spot, variability in leaf and fruit resistance among 17 varieties and 12 selections indicates feasibility of breeding for resistance, 112
---: limb dieback, perithecial stage of Eutypa armeniacae is only important source of field infection, 53
---: ring pox virus, 91 foreign varieties found to be hosts, 289
Arceuthobium campylopodum f. tsugensis: 921
Archtostaphylos manzanita: 181
--- viscida: 181
Arizona: 49; 145; 414; 512; 629
Arkansas: 862
Armillaria mellea: associated with dieback of Ceanothus spp. in Pacific Northwest, 399
Artemisia tridnetata: 181
Aspergillus niger: 175; causes new sorghum stalk rot, 352
Australia: 323; 494
Azalea: Rhizoctonia leaf blight, report on disease and control, 354

Bacteria: highly virulent Erwinia sp. was isolated from naturally infected lettuce and rapini plants and potato plants and tubers in Arizona, 414; method for bleaching leaves for microscope investigation of microflora on the leaf surface, 493; partial vacuum method of inoculating bean seeds with 3 common pathogens is rapid and efficient, and provides uniformly infected seed, 110; Pseudomonas glycinea, identification of 7 races on soybean, 557; serological relationships of races 1, 2, and 3 of Pseudomonas solanacearum, 275
Bacterial blight (see Blight)
--- canker: of tomato, serological techniques aid in detection of, 96
--- rot (see Rot)
--- spot: of apricot, variability in leaf and fruit resistance among apricots tested indicates feasibility of breeding for resistance, 112; of tomato, summary report over past 10 years on field evaluation and observations on seasonal occurrence, sources of primary inoculum, dissemination, and control in south Florida, 699
Bamboo: rust, first report of Uredo ignava in the United States, found in Florida, 112
Barley: loose smut, larger seeds produce drastically lower percentage of infected plants, 101
---: pathogenicity of Pythium aphanidermatum, 292
---: yellow dwarf virus, resistance in selected barley varieties, 380
Bean: asexual sporulation of Cephalosporium gregatum on autoclaved Golden Bantam sweet corn medium for culturing on Pinto 111 bean, 33; partial vacuum used to inoculate seeds with pathogenic bacteria, rapid and efficient method to supply uniformly infected seed, 110; survey of lima bean diseases in the South in 1965, 316
---: angular leaf spot associated with severe defoliation of red kidney beans in Pennsylvania, 494
---: bacterial blights: in Colorado, 10% loss in 1965, 371; inheritance of reaction to halo and common blight bacteria in Nebraska #1 sel. 27 dry bean line, 29; halo blight, quick laboratory test to determine pathogenicity of Pseudomonas phaseolicola, 238
---: capsule mold and capsule drop of castorbean, symptoms, losses, and varietal resistance, 35
---: downy mildew, effect of low post-inoculation temperature on colonization of lima bean stems by Phytophthora phaseoli, 734
---: powdery mildew: control by 2 new sulfites, 645; test of fungicides on snap beans, captan + tribasic copper sulfate gave excellent control without phytotoxicity, 847
---: root rot, identity and relative importance of major root rot pathogens of snap beans in Wisconsin, 578
---: soybean cyst nematode, reproduction on 12 hosts compared, 608
---: viruses: sources of virus resistance to bean yellow mosaic, alfalfa mosaic, western ringspot, and common bean mosaic in cultivars and breeding lines, 532; yellow mosaic virus, common pea mosaic virus strongly indicated as strain of, 337
Beet: (or tablebeet), evaluation tests in New York indicated that Rhizoctonia damping-off of seedlings was effectively controlled when PCNB was used in combination with other standard fungicides used to control the more important Pythium damping-off diseases, 614

Berry speckle: of cranberry, common but apparently undescribed, in Wisconsin, 539
Bioassay: possible systemic nature of six 8-quinolinol compounds tested by branch-injection into prune trees for Cytospora fungi control, 773
Biological control: in India the use of soil amendments may be practical substitute for expensive chemicals for lowering root-knot nematode populations in soil, 668; study of nematode-trapping fungi and predaceous nematodes occurring in Massachusetts cranberry soils with a view toward possible biological control of plant-parasitic nematodes, 584
Bipolaris sorokiniana-Fusarium nivale sequence in roots and crowns of winter wheat in Washington, 154
Birdsfoot trefoil: Leptodiscus terrestris, first report from Missouri as root-inhabitant fungus of a legume, 753; Thielaviopsis basicola development enhanced by temperature stress on roots in greenhouse-grown plants, 825
Blight: of Arizona cypress, new blight found in Georgia and South Carolina possibly caused by Monochaetia unicornis, 482; of guar, 9 fungicides tested for efficacy in control, 921; of maple, circumstantial evidence suggests that nematodes may contribute to incidence, 677
---: Alternaria, combined control of blight and powdery mildew of carrots, 594
---: bacterial: of beans in Colorado, 10% loss in 1965, 371; inheritance of reaction to halo and common blight bacteria in a bean cross, 29; of cotton, tests for races of Xanthomonas malvacearum from Australia yielded 13 infected varieties, 2 of which were resistant in the United States, 323; of radish leaf, highly pathogenic strain of Pseudomonas maculicola possibly resulted from repeated cropping on same ground, 192; of soybean, 16 methods tested for determining pathogenic races of pathogen, 560; of soybean, first report of identification of 7 pathogenic races, 557; Stemphylium and Xanthomonas blights, severity on tomato foliage affected by mode of irrigation in Israel, 635
---: boll, of flax, serious reduction of crop value in Texas, 840
---: brown spot needle, first report on jack pine, also found on lodgepole pine, Canada, 208
---: fire, of Stewart strain of Bartlett pear, further tests on susceptibility, 787
---: flower, of camellia, spread in Louisiana, 525
---: Fusarium, of turfgrasses, 942
---: gummy stem, of cantaloup, 1161 introductions screened for resistance, PI 140471 approaches immunity under field conditions, 661
---: halo: on bean, quick laboratory pathogenicity test, 238
---: leaf: of azalea, 354; northern leaf, factors affecting survival of Helminthosporium turcicum in corn leaf tissues, 377; Phytophthora leaf, of cacao seedlings, effect of adding DMSO to fungicides for control, 422
---: Rosellinia needle, on Douglas-fir seedlings, first report of R. herpotrichioides in California, 249
---: seedling, of cotton, use of deeply incorporated trifluralin herbicide predisposed plants to damping-off, PCNB for control, 172
---: southern: of peanut, PCNB effective control in irrigated fields, 855; of tomato, losses reduced by deep plowing, 443
Blossom blast: of cranberry, caused by Glomerella cingulata, resulted in reduced fruit set, 346
Botryosphaeria ribis: 122
Botryotinia ricini: 35
Botrytis allii: occurrence in yellow Gilboa onions which are being grown for seed production in Israel, 21
--- cinerea: testing thermal dusting with tetrachloroisophthalonitrile for control on geranium, 777
Breadfruit: a new lethal disease in the Pacific islands can kill all branches to the main trunk in 6 weeks on a robust, full bearing tree, 892
Bremia lactucae: 460
Broomrape: false, field observations on the response of Maryland tobacco to, first identified in southern Maryland in 1965, 627
Brown core and stem cavity browning: association on stored McIntosh apples, possibly both disorders have a common cause, 178
Brown patch disease: of St. Augustine grass, screening for resistance, 622
Brown rot: of oranges and grapefruit: in 3-year Israel experiments the rot-causing fungus was best con-

trolled when hot water was applied in groves 3 or 4 days after inoculation of the fruit, 867
Buckthorn: crown rust, 1000 isolates tested for pathogenicity on key differential oat varieties, 1961-1965, 935

Cacao: gall-inducing capacity of Fusarium rigidiuscula inactivated by interference from another Fusarium clone, 219; the roll of Fusarium roseum in inducing greenpoint cushion gall in the Caribbean and Latin America, 229; Phytophthora leaf blight, effect of adding DMSO to fungicides for control on seedlings, 422
California: 53; 168; 181; 182; 211; 245; 249; 283; 318; 327; 439; 479; 581; 768; 832
Calonectria rigidiuscula: 219
Camellia: flower blight, spread in Louisiana, 525
Canada: 62; 144; 208; 451; 695; 722; 750
Canker: a canker disease of red delicious apple in Guatemala, 694; previously undescribed branch and stem canker with severe resinosis on balsam fir in Anticosti Island, 722; sporophores of Polyporus tomentosus var. circinatus associated with basal cankers caused by fusiform rust fungus, 527
---: aspen, antibiotic tests for control on lodgepole and ponderosa pines, 741
---: Corynebacterium michiganense, diagnosis of the disease in tomato, 96
---: Cytospora, on honey locust, first report on thornless species, 13
---: European, development in California apple district, 182
---: pitch, fruiting of Fusarium lateritium pini on naturally infected pine, 476
Canna: mosaic disease caused by cucumber mosaic virus, 1st report of natural infection of canna by CMV in the United States, 899
Capsule mold and capsule drop: of castorbean, symptoms, losses, and varietal resistance, 35
Carrot: 681; combined control of Alternaria blight and powdery mildew, 594
Cassia: bioassay plant in method for detecting flexuous-rod virus in Cattleya leaf sap, 779
Castilleja miniata: uredial and telial stages of white spored Peridermium stalactiforme found on, Alberta, Canada, 144
Castor bean: influence of cropping to castor beans on populations of root-knot nematodes in soil, 132
Cattleya: infectious blossom necrosis, method for detecting flexuous-rod virus in leaf sap, 779
Ceanothus spp.: association of Armillaria mellea with severe dieback in Pacific Northwest, freeze of November 1955 may be predisposing factor, 399
Celery: ozone damage in Connecticut, 721
Cenangium singulare: 741
Cephalosporium spp.: 722; associated with a decline of oak in Texas, 75
--- gregatum: 598; asexual sporulation on autoclaved Golden Bantam sweet corn medium for culturing on Pinto 111 bean, 33
--- maydis: 401
--- stripe: of wheat, first published report of Cephalosporium gramineum in Michigan, 190
Ceratocystis fagacearum: 45; 736; 862
--- fimbriata: 741
--- ulmi: 473; 573; 618; 808
Cercis canadensis: 945
Cercospora aleuritidis: imperfect stage of angular leaf spot of tung, 54
--- beticola: 457
--- gigantea: 357
--- kikuchii: 910; cause of purple stain of soybean seedcoats, 464
Cercosporella herpotrichoides: 852
Chaetosiphon fragaefolii: 586
Chemicals: chemo- and thermotherapy for control of leaf curl virus of Malvaceous plants, 88; common names for pesticides, 725; effects of several on oak wilt organism in the field, 736; 50 compounds tested did not inactivate crinkle strawberry virus when sprayed on foliage of infected plants, 843; sensitivity tests with fungicides for control of cotton seedling disease pathogens, 301; soil treatments and plastic mulches affect development of new apple trees in many ways when used at planting time, 201; two new sulfite acaricides effective in controlling apple and bean powdery mildews, 645
---: creosote, urea, and liquid borate equally effective in preventing stump surface infection by Fomes annosus

after 10 months, 22
---: 2,3-dihydro-5-carboxanilido-6-methyl-1,4-oxathiin-4,4-dioxide, effective seed treatment for systemic control of stripe rust of wheat seedlings, 806
---: dimethyl sulfoxide, effect when added to fungicides for control of Phytophthora palmivora on cacao seedlings, 422
---: fluorescent Phosphor 2282 used to measure fungicide deposition and retention on watermelon foliage, 327
---: Freers' Dutch Elm Arrester (X-D.E.A.) tested for Dutch elm control, inconclusive results, 573
---: gibberellic acid sprays on sour cherry yellows virus infected trees increased fruit spur development at certain growth stages and environmental conditions, 240
---: growth regulators, effect of host growth inhibition by Cycocel on leaf rust infection types, 108
---: methionine and Dexon tested for control of pea root rot, divergence between results from field and greenhouse tests, 91
---: PCNB counteracts damping-off of cotton seedlings, which is a side effect of deeply incorporated trifluralin herbicide, 172
---: SMDC, DBCP, and DD used to control soil nematodes before isolation of soil fungi in a study of root rot of red pine seedling nursery, 545
---: tin compounds: fungicidal effect against 2 leaf diseases of carrot which so far could only be controlled by compounds of different chemical composition specific for each disease, 594
---: triphenyl tin combinations effective in control of tar spot of citrus, 357
---: urea-formaldehyde wet-strength resins caused lenticel injury to Delicious apples in storage in Washington, while polyamide resins did not, 885
Cherry (see also Prunus): Barbados-(Malpighia glabra): new spot anthracnose in India, possibly caused by Colletotrichum sp., 518
---: viruses: bud transmission of dwarfing virus, 367; severe fruit necrosis caused by strains of green ring mottle virus, 59; Shiro-fugen as indicator for Prunus ringspot virus complex, 278; sour cherry yellows virus, gibberellic acid sprays increased fruit spur development of infected trees at certain growth stages and environmental conditions, 240
Chile: 280
Chili (see Pepper)
Chokecherry: western X-disease virus widely distributed in North and South Dakota and Nebraska, threatens this shrub commonly used in windbreaks and as important component of wildlife plantings, 659
Chromatography: paper, of leaf tissue, direct method, 390
Chrysanthemum: wilt of, susceptibility of several cultivars to the Yellow Delaware Fusarium and effect of N nutrition on susceptibility, 882
Cicca acida (see Phyllanthus acidus)
Cissus rhombifolia: immune to Xiphinema index, 27
Citrus spp.: 681; storage life of oranges was extended by gamma radiation of fruit inoculated after picking with Penicillium digitatum and P. italicum, Israel, 874
---: dieback and gumming of 'Robinson' tangerine twigs caused by Diplodia natalensis in Florida, 429
---: nematodes: citrus-root nematode found in many sweet orange and sour lemon groves in Lebanon, 960; Tylenchulus semipenetrans (citrus nematode) found in Trinidad for first time, 478
---: Phytophthora citrophthora incubation period studied in inoculated trees under natural conditions, 251
---: postharvest decay: Miller's technique for impregnating wrappers with antifungal compounds, 782
---: rots: black core rot of mandarin oranges caused by Alternaria tenuis, 127; brown rot, hot water treatment for control on oranges and grapefruits is most effective when applied within 3 or 4 days after inoculation, Israel, 867; brown rot syndrome in Western Australian citrus orchards, 3 Phytophthora spp. associated with, 494; stem-end rot fungi in Florida lemons for fresh fruit market, 860
---: tar spot, chemical control, 357
---: viruses: Arizona's Cooperative Citrus Registration-Certification Program, organization, historical development, and progress, purpose = establishment of virus-free budwood, 145; psorosis virus, transmitted through seeds of at least one citrus hybrid ('Carrizo' citrange) at levels of 15 to 31%, 81

Cladosporium sp.: 35
--- fulvum: 686
Clathrus ruber: 840
Clover: subterranean, occurrence of wart disease in California believed to be first report on this host in the United States, 797
Club root: in combination with sugarbeet and cabbage root nematodes on Brussels sprouts, control by combination soil treatments, 133
Colletotrichum spp.: 313; 518
--- capsici: 299
--- --- f. cyamopsicola: 921
--- gloeosporioides: 385
--- graminicolum: of sorghum, spread in dead stalk tissue cells, and inhibition by living stalk tissue cells, 287
--- lagenarium: 340
--- lilii: 751
--- orbiculare: sexual expression in, 871
--- truncatum: 316
Colonization: of birdsfoot trefoil roots, by several fungal species, first report in Missouri for Leptodiscus terrestris, 753; intercellular, by 3 Malayan Fusarium isolates in Vigna and Hibiscus, 116
Colorado: 309; 371; 565
Common names: for pesticides, 725
Compatibility: 48 pesticide-nutrient mixtures compared in a 3-year field test on watermelon in Florida, 602
Conifers: distribution of Fomes annosus in natural forests of California, 832; 16 species inoculated with 7 isolates of Phytophthora cinnamomi from various hosts and locations provides evidence for existence of biotypes of the fungus, 647
Connecticut: 721
Coriander: new host for Erwinia aroideae (soft rot), 701
Corn: diseases in Kentucky in 1965, 218; sterilized Golden Bantam sweet, medium for asexual sporulation of Cephalosporium gregatum for culturing on Pinto 111 bean, 33
---: ergot conidia collected from sorghum will infect corn, 828
---: northern leaf blight, factors affecting survival of Helminthosporium turcicum in leaf tissues, 377
---: rots: inoculation and rating procedures for resistance of corn to stalk rot under conditions in the South, 655; reaction of corn inbred lines, varieties, and hybrids to 4 fungi causing stalk rots in Egypt, 401
---: rust, morphological evaluation of 2 pathogenic clonal lines of Puccinia sorghi and their inbred recombinants on a homozygous susceptible corn, 119
---: smuts: increase in susceptibility of inbred W22 by adding potassium to soil associated with decrease in reducing sugar and increase in nitrogen in the pith tissue, 12; head smut in Idaho, tests indicate fungus is not seedborne, 215
---: viruses: distribution and identification of mechanically transmissible viruses in Louisiana, 325; evidence that corn stunt virus may not occur in California, 318; occurrence of mechanically transmissible virus of corn in Mississippi, 409; sugarcane mosaic virus, reaction of 180 sweet and dent varieties to Johnson grass strain of, 793
---: ---: maize dwarf mosaic: history and distribution in Tennessee, 427; host range, properties, and partial purification of a Missouri isolate, 862; found in Iowa for first time, 88; presence established in several counties in Pennsylvania in 1965, 6; a heavy-duty mobile air blast inoculator has been designed, built, tested, and proved to be rapid and efficient for both field and greenhouse use, 363
---: ---: wheat streak mosaic: host range and spread of virus and vector in South Dakota, 1962-66, 819; tests indicate that Ohio 3A virus culture is strain of wheat streak mosaic virus, occurrence in Ohio suggests that wheat and corn could be year-round hosts for virus and wheat curl mites throughout corn belt and eastern Great Plains, 953
Corrections: 280; 536; 624
Corticium fuciforme: 525
Corynebacterium flaccumfaciens: 110
--- michiganense: 96
Cotton: reduction in yield caused by diseases in 1965, 350
---: angular leaf spot, eradicated from California through sanitation, 168
---: bacterial blight, tests for races of Xanthomonas malvacearum from Australia yielded 13 infected varieties, 2 of which were resistant in the United States, 323
---: boll rots, method for estimating losses, 254
---: nematodes: plant-parasitic nematodes, seriousness and scope of infestation in Lower Rio Grande Valley of Texas, 149; root-knot

nematode, testing experimental nematocides on 'Deltapine Smoothleaf' under Arizona conditions, 512
---: Rhizoctonia solani isolated from seed that were acid delinted and surface sterilized with alcoholic mercuric-chloride, 770
---: seedling damping-off, a side effect of deeply incorporated trifluralin herbicide counteracted by PCNB, 172
---: seedling disease complex variability in New Mexico, in relation to chemical control, 301
---: Verticillium wilt, comparison of virulence of isolates of Verticillium albo-atrum from several cotton-growing regions in the United States, Mexico and Peru, 930
Cranberry: application of pesticides with "ground effect" machine, 715; study of nematophagous fungi and predaceous nematodes associated with cranberry soils in Massachusetts, 584
---: berry speckle, in Wisconsin, evaluation of fungicides and spray schedules for control, 539
---: Glomerella cingulata infection associated with increased blossom blast and decreased fruit set; fruit rot symptom expression also dependent upon high or low N, P, K, or S levels, 346
---: plant-parasitic nematodes, in Wisconsin, 957
---: ringspot disease, virus-like ringspot symptoms on fruits and leaves of Searles and Howes varieties in Wisconsin, 543
Crataegus oxyacantha: Gymnosporangium confusum on, in California, 768
Crater canker: of Golden Delicious apple, bark disease of unknown cause, extent of tree decline is proportional to amount of trunk injury, 3
Cronartium comandrae: 528
--- comptoniae: 419; 744
--- fusiforme: 388; 527; development of pycnia and aecia in albino form on inoculated pines, 842
--- ribicola: 554
Crucifers: new hosts for Meloidogyne hapla, 523
---: bacterial leaf blight: of radish, highly pathogenic strain of Pseudomonas maculicola possibly resulted from repeated cropping on same ground, 192
---: cabbage leaf spot, transmission of by slugs in France, 764
---: nematodes: sugarbeet and cabbage root nematodes occurring with club root organism on Brussels sprouts, marked yield increase with control, 133; sugarbeet nematodes associated with reduced yields and quality in cauliflower, cabbage, broccoli, Swiss chard and mustard, 611
Cucumber (see Cucurbits)
Cucurbits: decay of melons from California and Mexico on the New York market, caused by Geotrichum candidum, 222; viruses of, on the west coast of Mexico, 947
---: cantaloup: gummy stem blight, 1161 introductions screened for resistance, PI 140471 approaches immunity under field conditions, 661
---: cucumber: 681; anthracnose, downy mildew, and soil rot, control, 340
---: muskmelon necrotic fleck virus, new strain of tobacco mosaic virus isolated from muskmelon in Arizona, 49
---: squash: 681
---: watermelon: 3-year compatibility field tests among fungicide-insecticide-nutrient sprays in Florida, 602; surfactant influence on deposition and retention of fungicides on foliage, 327; mosaic virus, aluminum foil tested for repellency to melon aphids, 305
Cultures: of Cephalosporium gregatum on Pinto 111 bean, autoclaved Golden Bantam sweet corn medium for asexual sporulation of C. gregatum, 33; of 4 Geortrichum candidum isolates on four agars in a study of plant tissue pH and susceptibility to plant and animal isolates, 681; induced sporulation of fungi on non-sterile filter paper in a moist chamber, 789; simple and inexpensive device for uniform transfer of fungus inoculum, 345; preservation of, 32 Helminthosporium spp. were stored under mineral oil on agar slants, 321; survival of various smut fungi cultures for 11 years under mineral oil, 800; sexual expression in Colletotrichum orbiculare, 871; in vitro increase in virus infectivity, 639
Curvularia geniculata: first report on Sambucus species, found on S. chinensis, 488
--- verruculosa: causes severe disease of rice in India, 314
Cuscuta cassytoides: 593
Cylindrocarpon sp.: 722

Cypress: Arizona, new blight found in Georgia and South Carolina possibly caused by Monochaetia unicornis, 482
Cytospora spp.: 773
--- chrysosperma: 741
--- gleditschiae: 13
Cytosporina sp.: perithecial stage of Eutypa armeniacae is only important source of apricot limb dieback field infection, not this imperfect stage, 53

Dalbulus maidis: 427; 570
Damping-off: of alfalfa, caused by Xanthomonas alfalfae, in Kansas, 731; of alfalfa seedlings, effects of 4 fungicides on plants and on control of disease in sand, 435; detection of Pythium in soil, 791; of table beet, controlled by combination fungicides for Rhizoctonia and Pythium organisms, 614; of Tephrosia vogelii, control studies with fungicides in Puerto Rico, 484; of tomato, 124
Decay: of western melons on the New York market, caused by Geotrichum candidum, 222
Decline: of mature apple trees associated with plant-parasitic nematodes, effect of DBCP and cynem, 877; of loblolly pine, associated with Phytophthora cinnamomi and Pythium spp., 596; of oak in Texas, associated with Cephalosporium, 75
Dieback: apricot limb, perithecial stage of Eutypa armeniaceae is only important source of field infection, 53; of Ceanothus spp., association of Armillaria mellea with, 399; and gumming of 'Robinson' tangerine twigs caused by Diplodia natalensis in Florida, 429; of maple, circumstantial evidence suggests that nematodes may contribute to incidence, 677; and mortality, of red and jack pines caused by Scleroderris lagerbergii in Upper Michigan, 402
Dieffenbachia spp.: brown leaf spot, severe disease in Florida, proved to be caused by Leptosphaeria sp., 687
Dimpling: of McIntosh and Cortland apples, possibly physiological disease, 151
Dioscorea alata: anthracnose of, efficacy of different fungicides for control, 385
Diplodia natalensis: 429; 860
--- zeae: 655
Diseases: check list of plant diseases previously unreported in Hawaii, 411; in Kentucky in 1965, 218; of lima beans in the South in 1965, 316; reduction in yield of cotton caused by, 350
Distribution: amendments to records of Cronartium comptoniae, 744
Dodder: parasitic on 2 principal and 3 secondary hardwood hosts each in a different family, Tanzania, 593; virus resembling tobacco streak isolated from dodder growing on a desert shrub in southern California, shrub may be reservoir for virus, 479
Dothidella ulei: five clones of Hevea sp. designated as differentials for identifying races of, 187
Dothistroma pini: on Pinus radiata, first report of this fungus in Chile, 280
Downy mildew: of cucumber, control, 340; of lettuce, immune varieties Valverde, Calamar and Valmaine seriously affected by new race of Bremia lactucae in south Texas, 460; of lima bean, effect of low postinoculation temperature on colonization of stems by Phytophthora phaseoli, 734; of peas in western Europe, fungicidal experiments indicate that 1 spraying at the onset of flowering can reduce number of diseased pods by about half, 257; of sorghum, new sources of resistance, 641
Drought: artificially created on loblolly pine stumps increased penetration of stumps by Fomes annosus 3 months after inoculation, 747
Dutch elm disease: control studies, method for inoculation, 808; effects of SMDC on elm roots, 473; further tests with various antibiotics and fungicides gave inconclusive results; however, captan + DMSO was the best of those tested, 573; known distribution in Nebraska, 1965, 618
Dwarfmistletoe: found on Sitka spruce in Alaska, new host record, 921

Egypt: 401
Elm: Dutch: effects of SMDC on elm roots, concentrations, time, and control tested, 473; further tests with various antibiotics and fungicides gave inconclusive results; however, captan + DMSO showed promise, 573; known distribution in Nebraska, 1965, 618; method for inoculation of trees with spore suspension of Ceratocystis ulmi for

control studies, 808
---: phloem necrosis virus (Morsus ulmi) of American elm in Mississippi Delta, 751
Enzymes: pectic, production of by Malayan isolates of Fusarium, 116
Epidemiology: of bacterial spot of tomato, a study in south Florida of seasonal occurrence, sources of primary inoculum, dissemination and control, 699
Eradication: of Xanthomonas malvacearum from California through sanitation, 168
Ergot: of sorghum, host range and epidemiology in Nigeria, 828
Erwinia sp.: highly virulent isolate from Arizona vegetables, 414
--- amylovora: 787
--- aroideae: 701
--- carotovora: 938
Erysiphe polygoni: 645; 847
--- umbelliferarum: 594
European canker: development in a California apple district, importance of ascospores and macroconidia, susceptibility of leaf scars, and of rain, 182
Eutypa armeniacae: perfect stage recognized as cause of apricot limb dieback in California, not imperfect stage (Cytosporina), 53

Fertilizer: application of potassium to soil greatly increased smut in corn inbred W22 by decreasing reducing sugar and increasing nitrogen in the pith tissue, 12
Fir (see Abies and Pseudotsuga)
Flax: rust, 16 races in varying frequencies in Canadian prairies and in North Dakota in 1963 and 1964, 62; rust, pasmo, and boll blight diseases and weathering of seed and fiber production accounted for 75 to 80% reduction in potential value of crop in Texas during 1965-66 growing season, 840
Florida: 112; 124; 292; 602; 687; 689; 707; 822; 860
Fomes annosus: 446; 589; 747; development of a hazard rating for, in pine stands, 510; distribution in natural forests of California, 832; found attacking and killing five previously unreported hosts, California, 181; root disease survey to determine extent of damage in natural forest stand shows that fungus is greater menace in California than past reports indicate, 439; three stump treatment chemicals effectively prevented infection by, 22
Fraxinus americana: graft transmission of a ringspot-like virus from, 905
Frost: injury to wheat, 928
Fungi: associated with wilts and root rots of alfalfa in eastern Ontario, and an assess of damage, 695; evaluation of soil microflora and isolation by apple technique in relation to the root rot disease of red pine, 545; 11 forest fungi tested for effects of cycloheximide in culture, 497; induced sporulation on filter paper, 789; inoculation technique using nylon mesh leaves fungus free of adhering growth media, can be adapted to many fungus-host combinations, 122; isolated from hypocotyls and cotyledons of fungicide-treated peanut seed grown in infested soil in Georgia, 162; Nematophagous, studied in cranberry soils in Massachusetts as possible biological control agents of phytoparasitic nematodes, 584; new records of powdery mildew fungi, 709; pepper seed microflora inhibition by antibiotic, Pentaene G8, 299; a simple and inexpensive device developed for "one-step" aseptic uniform transfer of fungus inoculum, 345; study of the important root-inhabiting fungi associated with birdsfoot trefoil in Missouri, 753; stinkhorn, possibly Clathrus ruber, unusual occurrence on golf turf, 840
Fungicide: effect of treatment of peanut seed on soil-borne fungi at the hypocotyl and cotyledonary surfaces, 162; evaluation of fungicides and spray schedules for control of berry speckle disease of cranberry, 539; four new formulations tested for control of pecan scab for 3 seasons in Oklahoma, 599; 3-year field test of compatibility with insecticide-nutrient sprays for disease and insect control on watermelon, 602; Miller's technique for impregnating wrappers with antifungal compounds, 782; nine tested for efficacy in controlling blight of guar, 921; several evaluated for control of angular leaf spot of tung, 717
---: carbamate and copper fungicides tested for efficacy in controlling anthracnose of Dioscorea alata in India, 385
---: new carbamoyloxime insecticide (UC 21149, Temik) shows systemic fungicidal activity in laboratory

studies, 424
---: cycloheximide, effects on 11 selected forest fungi in culture, 497
---: DCNA, for control of Rhizopus rot on peaches, 565
---: maneb, one spraying with maneb or a a coordination product of zinc ion and maneb at onset of flowering can reduce number of downy mildew infected pea pods by about 50% in western Europe, 257
---: 1,4-oxathiin, 2 derivatives show chemotherapeutic activity against smut and rust diseases of grasses, 624
---: pentachloronitrobenzene: effective in controlling Sclerotium rolfsii when southern blight is principal disease problem in irrigated peanuts, 855; PCNB added to currently recommended table beet seed ball treatment fungicides results in the control of Rhizoctonia solani as well as of 2 Pythium species which are primarily responsible for damping-off of seedlings in New York, 614
---: Pentaene G8, see antibiotics, 299
---: several tested together: effect of various combinations of 8 on stand establishment, maintenance, and rotenoid concentration of Tephrosia vogelii in Puerto Rico, 484; captan + dimethyl sulfoxide was only material that showed promise of several antifungal antibiotics and fungicides tested for Dutch elm disease control, 573; captan + tribasic copper sulfate (one of several tested) gave excellent control of powdery mildew on snap beans without phytotoxicity, 847; neutral copper, agricultural streptomycin and maneb effective against bacterial spot disease of tomato in south Florida, 699; cycloheximide and phytoactin, evaluation of basal stem application of, for control of white pine blister rust on sugar pine, 554; Daconil, Dithane M-22, Dithane M-22 special, Dithane M-45, Manzate, and Manzate-D compared for control of anthracnose and downy mildew of cucumbers, 340; effect of adding DMSO to 3 copper fungicides and to malachite green for control of Phytophthora palmivora leaf blight on cacao seedlings, 422; nickel + dithiocarbamates, evaluation for control of wheat stem rust, 373; bioassay with six 8-quinolinol compounds for control of Cytospora fungi in prune trees indicates some promise of effective systemic action, 773; PCNB, Thimer, and thiram tested for control of snow mold on four varieties of bluegrass in Alaska, 69; combination with surfactants, retention of on watermelon foliage measured by a fluorescent chemical, 327; effects of thiram, captan, pentachloronitrobenzene and chloranil on alfalfa seedlings and on controlling Rhizoctonia solani in sand, 435
---: SMDC, effects on elm roots, concentration-time-control tests on Ceratocystis ulmi, 473
---: sodium-o-phenylphenate, 2-aminobutane, and thiabendazole tested for control of blue-mold rot of apples, 913
---: tetrachloroisophthalonitrile, thermal dusting for control of Botrytis cinerea on geranium in the greenhouse, 777
Fusarium spp.: 122; 695; colonization by 3 Malayan isolates in Vigna and Hibiscus, 116; toxin production by non wilt-inducing isolates, 261
--- decemcellulare: 229
--- graminearum: 401
--- lateritium pini: fruiting on naturally infected pine pitch cankers in North Carolina, 476
--- moniliforme: 401; 655
--- nivale: 154
--- oxysporum: 469; 882
--- --- f. apii: 391
--- --- f. lycopersici: 707; serological differentiation of race 1 from race 2, 444
--- rigidiuscula: 219
--- roseum: 26; 229; 942
--- solani: 160; pathogenic to bean and pea, 578
--- wilt (see Wilt, Fusarium)
Fusicladium effusum: 599

Ganoderma lucidum: 243
Georgia: 162; 306; 393; 482; 839
Geotrichum candidum: 222; 3 plant and 1 human isolate of: study to test pathogenicity of and relationship between tissue pH and susceptibility of 15 puncture-wound inoculated plant tissues, 681
Geranium: bacterial stem rot and leaf spot, satisfactory technique for checking presence of bacteria in commercial plants, 515; Botrytis cinerea, testing thermal dusting with tetrachloroisophthalonitrile for control, 777
Gibbera compacta: 539

Gloeosporium perennans: 348
Glomerella cingulata: 346
Gloxinia: Thielaviopsis basicola rot of rhizomes from Europe, 709
Gramineae: pathogenicity of 91 morphologically similar isolates of Helminthosporium to 30 gramineous species, 382
Grape: hot-water treatments effective in eliminating Pratylenchus vulnus from grapevine roots, however soil treatments should still be used, 858
Grasses: citrus nematode found on an indigenous Florida grass, Andropogon rhizomatus, 822; Hypsoperine graminis in Kansas, 396; root damage resulting from HCN produced by Marasmius oreades, 264; 2 new systemic fungicides (derivatives of 1,4-oxathiin) show chemotherapeutic activity against rusts and smuts, 624; 44 tested for susceptibility to aster yellows virus, host range now includes 8 grasses (including oats), 17
---: Bermuda grass, new host for red thread, found in Mississippi, 525
---: bluegrass, snow mold of, three chemicals tested for control on four varieties of bluegrass in Alaska, 69
---: gamagrass: new host of Dalbulus maidis, 3rd plant species reported to be accepted by leafhopper for development from egg to adult, 570; new perennial host of maize dwarf mosaic virus, Illinois, 65
---: guinea, host range and epidemiology of the sorghum ergot organism in Nigeria, 828
---: Johnson grass: maize dwarf mosaic virus, positive transmission from one source, first report in Kansas in 1965, 11; 2 distinct strains of maize dwarf mosaic virus in northeastern U.S. identified by host reaction and serological relationships, 814
---: Kentucky bluegrass: evidence for existence of physiologic races of stripe smut, 430; severity of melting-out reduced by use of resistant varieties, avoiding excessive nitrogen fertilization, and raising cutting height of turf, 703
---: St. Augustine: screening for tolerance to Rhizoctonia solani, 622; seedlings mechanically inoculated with maize dwarf mosaic virus resulted in 80% infection, potential threat to South, 441
---: turfgrass: effects of nematocides on growth, 489; Fusarium blight of turfgrasses, 942; pseudo-root-knot nematode, found in Texas for first time, 645; unusual occurrence of stinkhorn fungus on golf turf, has unsightly appearance, viscous nature, offensive odor, and distorts putting surface, 840
---: turf and forage, studies of stripe smut prevalence, pathogenicity and control in New Jersey, 294
---: western wheatgrass, first record of seed-gall nematode on, found in Montana, 45
---: zoysia: first report of rust on Zoysia spp. in Alabama, 885; occurrence of rust in Georgia, 839; Puccinia zoysiae found on Zoysia spp. in Mississippi, 770
Greenpoint cushion gall: of cacao, the role of Fusarium roseum in causing the disease, 229; of cacao, gall-inducing capacity of Fusarium rigidiuscula inactivated by interference from another Fusarium clone, 219
Guar: blight of, 9 fungicides tested for efficacy in control, 921; tobacco ring spot virus, new occurrences in Texas and Oklahoma, 7
Guatemala: 694
Guignardia vaccinii: 539
Gymnosporangium confusum: in California, first report in western hemisphere, 768

Hardwoods: dodder parasitic on seedlings in Tanzania forest nursery, 593
Hawaii: 411; 519
Hazard rating: for Fomes annosus in Virginia, guidelines provided by a survey which are based on criteria of soil texture, depth of A horizon and height of water table to predict rate of spread in infected loblolly pine stands, 510
Helminthosporium spp.: longevity and pathogenicity of 32 species stored under mineral oil on agar slants, 321; pathogenicity of 91 morphologically similar isolates to 30 gramineous species, 382
--- turcicum: factors affecting survival in corn leaf tissues, 377
--- vagans: 703
Hemicycliophora arenaria: 581
Hemp sesbania: comparative rates of reproduction of soybean cyst nematode on 12 hosts, 608
Herbicides: diuron, fall applications to control weeds in winter wheat reduced incidence and severity of foot rot, 852

---: trifluralin predisposed cotton to Rhizoctonia seedling blight, PCNB may be practical control, 172
Heterodera cruciferae: 133
--- glycines: 15; 548; 608
--- punctata: occurrence and distribution, 869
--- schachtii: 133; 650
--- tabacum: 506
Hibiscus esculentus: effect of culture filtrates of Fusarium isolates on wilt production, 261; intercellular colonization and the production of pectic enzymes by Malayan isolates of Fusarium, 116
--- rosa-sinensis: leaf curl virus, chemo- and thermotherapy for control, 88
Honeylocust: Cytospora canker on, first report on thornless species, 13; Polyporus lucidus associated with sudden death of tree in Mississippi, 243
Horizon, A: depth of, as it relates to loblolly pine infection by Fomes annosus, 510
Hosts: 10 weed species collected in Mississippi suspected to be natural summer hosts for barley yellow dwarf virus, 692
Hot water treatments: (see temperature)
Hydrogen cyanide: production by Marasmium oreades damages cereal and grass roots, 264
Hymenoclea salsola: serves as native host to sheath nematode which is a native to California soils, 581
Hypoxylon pruinatum: 741
Hypsoperine graminis: 645; in Kansas, 396

Idaho: 215
Ilex spp.: 4 cultivars tested for pathogenicity of Meloidogyne spp., 664
Illinois: 65; 446
India: 313; 314; 612; 668; 701
Insecticides: combined with fungicides and nutrients in 3-year compatibility tests for spraying watermelon foliage in Florida, 602
---: 2-methyl-2-(methylthio)propionaldehyde O-(methylcarbamoyl)oxime, a new insecticide-acaricide, shows nematocidal and systemic fungicidal properties in the laboratory, 424
International facility: to prevent importation of foreign plant pathogens, 803
Interregional Research Project-2 (IR-2): Deciduous Tree Fruit Virus Project, originally designed to index viruses of stone fruits, has been extended to include indexing for latent viruses of pome fruits, particularly apples, 461
Iowa: 88
Irrigation: relationship between mode of irrigation and severity of tomato foliage diseases in Israel, 635
Isariopsis griseola: 494
Isolation: quarantine laboratory, preliminary outline of requirements needed by such a research facility (e.g. security to prevent escape) for the study of plant pathogens, viruses in particular, that have so far been excluded from entry into various countries, 803
Israel: 21; 711; 867; 874

Juniperus sabina: Gymnosporangium confusum on, in California, 768

Kansas: 11; 234; 396; 731
Kentucky: 218

Leaf: bleaching of, for microscope investigation of microflora on leaf surface, 493; disposable leaf presses for obtaining plant sap for serodiagnosis, 822
Leaf press: disposable, to obtain sap from plants for serodiagnosis, 822
Leaf scald: of sugarcane, found in Puerto Rico, first report of occurrence in one of the larger islands of the Antilles, 569
Leaf spot: angular: associated with severe defoliation of red kidney beans in Pennsylvania, 494; of cotton, Xanthomonas malvacearum eradicated from California through sanitation practices, 168; of strawberry, found in Wisconsin for first time, 167; of tung, effect on yield and oil content of fruit from trees sprayed with Bordeaux mixture or petroleum oil for 5 years, 54; of tung, evaluation of various fungicides for control, 717
---: bacterial: of alfalfa and new hosts, distribution in Kansas, 731; of geranium, inoculation technique, 515
---: brown, severe on several Dieffenbachia spp., proved to be caused by Leptosphaeria sp., 687
---: of cabbage, transmitted by slugs, 764
---: Cercospora, of peanut, technique for forecasting, 810
---: of strawberry, relative resistance of 63 varieties and selections studied at 2 locations for 2 years, 105

Lebanon: 960
Lecanosticta acicola: 208
Lenticel injury: of apples in storage caused by urea-formaldehyde resins, extent of injury also dependent upon temperature in Washington, 885
Leptodiscus terrestris: colonization of birdsfoot trefoil roots in Missouri, first report of, 753
Leptosphaeria sp.: proved to cause brown leaf spot of Dieffenbachia, 687
Lespedeza: comparative rates of reproduction of soybean cyst nematode on 12 hosts, 608
Lettuce: bacterial soft rot of Iceberg (Great Lakes) in Florida Everglades, 938
---: calico, caused by alfalfa mosaic virus not by tobacco ringspot virus, 629
---: downy mildew, immune varieties Valverde, Calamar and Valmaine seriously affected by new physiologic race of Bremia lactucae in south Texas, 460
---: highly virulent Erwinia sp. isolated from naturally infected plants in Arizona, 414
Libocedrus decurrens: Fomes annosus found killing trees within native range in California, 181
Lily: Easter: anthracnose on blossoms caused by black-scale Colletotrichum, 751; root-lesion nematodes, control with phosphate pesticides, 923
---: Lilium regale, bulbs from the Netherlands infected with Aphelenchoides fragariae were intercepted by plant quarantine inspectors, 798
Longidorus vineacola: 711
Loose smut: in Funza barley, influence of seed size on incidence of disease, 101
Lophophacidium hyperboreum: pathogenicity of, 356
Losses: tomato bacterial spot, evaluation of causes of this important disease and effective methods to combat its development and dissemination in south Florida, 699
Louisiana: 325; 525; 596; 686

Macrophomina phaseoli: 352; found on 3 new hosts in Pakistan, 171; responsible for most losses in forest nurseries in California, control by soil fumigation, 245
Macrosteles fascifrons: 17
Malvaviscus arboreus: leaf curl virus, chemo- and thermotherapy for control, 88
Marasmius oreades: fairy ring fungus produces HCN which damages grass roots, 264
Maryland: 351; 627
Massachusetts: 151; 584
Melampsora lini: 62; 840
Meloidogyne spp.: 126; 149; 664; 915
--- hapla: 132; new hosts of, 523
--- incognita: 132
--- --- acrita: 512
--- javanica: 132; 668
--- thamesii: 132
Melting-out: of Kentucky bluegrass, severity reduced by use of resistant varieties, avoiding excessive nitrogen fertilization, and raising mowing height of turf, 703
Methods and techniques: aluminum foil tested for repellency to melon aphids in watermelon mosaic virus control, 305; apparatus for mist culture of wheat, for study of influence of stripe rust on root development, 369; apple technique employed to isolate fungi from red pine seedling bed nursery in West Va. previously treated with several nematocides, 545; bleaching leaves for microscope investigation of microflora on the leaf surface, 493; comparison of 2 methods for inoculating Nuttall oaks with heart rot fungi, 552; deep plowing reduces losses due to southern blight on tomato, and generally improves plant vigor and fruit yields, 443; detection of Pythium in soil, 791; direct method for qualitative paper chromatography of leaf tissue, 390; disposable leaf presses for obtaining plant sap for serodiagnosis, 822; effects of surfactants on deposition and retention of fungicides on watermelon foliage measured by a fluorescent chemical, 327; estimating losses from cotton boll rots, 254; for evaluating hundreds of lines and varieties of wheat at maturity for resistance to common root rot, 26; forecasting leafspot on peanuts, 810; "ground effect" machine shows promise as a vehicle for applying pesticides on cranberry, 715; a heavy-duty, rapid, mobile and efficient pressure inoculator for mechanical transmission of maize dwarf mosaic virus was designed, built, and tested for both greenhouse and field use, 363; increment core sampling of Pinus spp. for Fomes annosus, 589; for identification of Puccinia graminis, permits sanitation, positive separa-

tion, and reduction in time needed to identify components of racial mixtures, 312; induced sporulation of fungi on filter paper, simple and inexpensive method, 789; inoculating geranium cuttings with Xanthomonas pelargonii to determine presence of bacteria in commercial plants, 515; inoculation of elms with spore suspension of Ceratocystis ulmi, 808; inoculation method is most efficient and precise of 16 tested for determining pathogenic races of soybean bacterial blight pathogen, 560; inoculation technique using nylon mesh can be adapted to many fungus-host combinations, 122; in vitro increase in virus infectivity, 639; isolating Verticillium spp. from field soil, 897; metal leaf presses in pliers to express plant sap for serodiagnosis, 822; Miller's technique for impregnating wrappers with antifungal compounds, 782; needle weight determination method accurately estimates intensity of Rhabdocline needle cast of Douglas-fir, 406; partial vacuum method of inoculating bean seeds with pathogenic bacteria is rapid and efficient, and provides uniformly infected seed, 110; quick laboratory test to determine pathogenicity of Pseudomonas phaseolicola on bean, 238; rapid flotation-sieving technique for extracting nematodes from soil, does not vary with time of year, 954; rapid method of assaying pathogenic potential of Puccinia graminis tritici populations for use in epidemic prediction and in detection of new, dangerous virulences, 673; in rating procedures for evaluating corn resistance to stalk rot caused by Diplodia zeae and Fusarium moniliforme, the toothpick and syringe methods were superior to inoculator method in the South, 655; screening St. Augustine grass for tolerance to Rhizoctonia solani, 622; storage of pycniospores of Puccinia graminis secalis in liquid nitrogen, 576; serological procedures simplify differentiation of race 1 from race 2 of Fusarium oxysporum f. sp. lycopersici, 444; simple device for uniform transfer of fungus inoculum, 345; a simple, efficient, and inexpensive method of collecting strawberry leaves for virus infection indexing, 426; simple, inexpensive, continuous sampling spore trap, 906; simple method for detection of a flexuous-rod virus associated with infectious Cattleya blossom necrosis, 779; storage of urediospores of Puccinia graminis tritici in liquid nitrogen, 502; survival of smut cultures under mineral oil for 11 years, 800; two methods for test inoculation of ponderosa pine with western gall rust, 550

Mexico: 598; 947

Michigan: 190; 402

Millet: ergot conidia collected from guinea grass will infect millet, 828

Minnesota: 557

Mississippi: 35; 243; 388; 409; 525; 570; 692; 770

Missouri: 753; 862

Mock orange: powdery mildew found for first time in Southeast (Georgia), 306

Monochaetia unicornis: 482

Montana: 45

Morus tatarica: inoculation technique to test pathogenicity of Fusarium spp. on, 122

Muskmelon (see Cucurbits)

Mycosphaerella aleuritidis: 717; perfect stage of angular leaf spot of tung, 54

--- citrullina: 661

--- fragariae: 105

Myrica gale: sweet fern blister rust on, 2 specimens found in Northwest Territories, 419

Myzus persicae: 613

Nebraska: 618

Nectria galligena: 182

Needle cast: Rhabdocline, of Douglas-fir, needle weight determination method accurately estimates intensity of disease, 406

Nematocides: effects on turfgrass growth, 489; new carbamoyloxime insecticide (UC 21149, Temik) shows nematocidal activity in laboratory studies, 424; phosphate pesticides control root-lesion nematodes in Oregon Easter lily plantings, 923; volatile and nonvolatile tested for control of tobacco cyst nematode, 507

---: 1,3-D and DD mixture, tested over a 2-year period for control of sugarbeet nematode in field plots cropped to vegetables, 611

---: DBCP: more effective than cynem when response of mature apple trees to both treatments was tested in the Hudson Valley, 877; 1,2-dibromo-3-chloropropane, 1,3-dichloropropene, and 1,2-dibromoethane tested for effect of exposure time and dose

on nematode eradication, under controlled conditions, 915
---: O,O-diethyl O-2-pyrazinyl phosphorothioate, used as a dip eradicated soybean cyst nematodes from roots of plants in pot tests, 548
Nematodes: in maple blight and dieback areas in Wisconsin may contribute to incidence of these diseases, 677; rapid flotation-sieving technique for extracting nematodes from soil, does not vary with time of year, 954; relation between nematocide soil treatment for control in red pine seedling bed nursery and effect on soil fungi isolated in study on cause of root rot of conifers, 545; survey of apple orchards in Hudson Valley, 1956-65, 783
---: Aphelenchoides fragariae, found on 3 tree peony species from Japan and on Lilium regale bulbs from the Netherlands by U.S. plant quarantine inspectors, 798
---: citrus: found on new grass host and associated with additional plants in two surveys, 822; found in Trinidad for first time, 478
---: citrus-root, in many sweet orange and sour lemon groves in Lebanon, 960
---: cotton root-knot, efficacy and residual nature of experimental nematocides tested for control on Deltapine cotton in Arizona, 512
---: cyst, notes on occurrence and distribution of Heterodera punctata, 869
---: dagger, can cause severe decline of Vinca minor, 420
---: ectoparasitic, disease of onion associated with Longidorus vineacola serious in Israel, 711
---: foliar, cause of "leaf stripe" disease of rubber plant in south Florida, 689
---: parasitic, effects of nematocides on turfgrass growth, 489
---: plant-parasitic: of cotton, percentage of infestation in Lower Rio Grande Valley of Texas, 149; on cranberries in Wisconsin, 957; response of mature apple trees to DBCP and cynem in the Hudson Valley, 877
---: predaceous, extracted from cranberry soils, identified and frequency counts made for future application to biological control of plant-parasitic species, 584
---: pseudo-root-knot, of turf grasses, found in Texas for first time, 645
---: reniform, varietal reaction of 24 sweetpotato selections to, 500
---: root-knot: effects of exposure time and nematocide dose on eradication, under controlled conditions, 915; pathogenicity of Meloidogyne spp. on Ilex cultivars, 664; Meloidogyne (may be undescribed species) possibly associated with decline of ponderosa pine in New Mexico, 126; new hosts for Meloidogyne hapla, 523; cropping to castor beans reduced Meloidogyne hapla and M. thamesii, but reproduction by M. incognita and M. javanica was excellent, 132; (Meloidogyne javanica) significant reduction of eggs, larvae, and mature females in soil as result of oil-cakes of margosa, castor, and peanut in India, 668
---: root-lesion: control in Oregon Easter lily plantings with phosphate pesticides, 923; hot-water treatments effective in eliminating Pratylenchus vulnus from grapevine roots, however soil treatment should still be used, 858
---: seed-gall, first record on western wheatgrass, and first record on grass in Montana, 45
---: sheath, a native to California soils, and Hymenoclea salsola serves as a native host, 581
---: soybean cyst nematode: additional weed hosts, 15; comparative rates of reproduction on 12 host plants, 608; elimination from roots of plants by chemical dips in pot tests, 548
---: sugarbeet: associated with reduced yields and quality in vegetable crops, 2-year field experiments with 1,3-D and DD mixture for control, 611; evaluating sugarbeet varieties and breeding lines for tolerance to nematode invasion, no outstanding resistance was observed, 650; possible interaction between sugarbeet and cabbage root nematodes and club root organism on Brussels sprouts, evaluation of combination soil treatments for control, 133
---: tobacco cyst, volatile and nonvolatile nematocides tested for control, 507
---: Xiphinema index, 5 new hosts for, 27
New diseases: blight of Arizona cypress found in Georgia and South Carolina possibly caused by Monochaetia unicornis, 482; new disorder of strawberry fruit, 327; Pingalap disease of breadfruit in the Pacific islands has destroyed an estimated 80,000 trees in 2 years, it can kill

all branches to the main trunk in 6 weeks, 892; previously undescribed branch and stem canker with severe resinosis on balsam fir in Canada, 722; sorghum stalk rot caused by Aspergillus niger, 352; stunt disease of peanut, undescribed virus causes severe dwarfing and fruit suppression, found in Virginia, 139
---: in India: new spot anthracnose of Malpighia glabra (Barbados-cherry) possibly caused by Colletotrichum sp., 518; anthracnose of Phyllanthus acidus, 313
---: in the United States: crater canker of Golden Delicious apple trees, bark disease of unknown cause, Washington, 3; ozone injury to celery in Connecticut is possibly first report in United States, 721
New distribution: check list of plant diseases previously unreported in Hawaii, 411
---: apple fruit rot, caused by Phialophora malorum, in Pennsylvania, 701
---: angular leaf spot of strawberry in Wisconsin, 167
---: camellia flower blight in southwest Louisiana, 525
---: Dothistroma pini in Chile, found on Pinus radiata, 280
---: false broomrape on tobacco in Maryland, 627
---: Fomes annosus found on Libocedrus decurrens in California, 181
---: Gymnosporangium confusum in western hemisphere for first time, found in California, 768
---: Lecanosticta acicola found in Canada, 208
---: Macrophomina phaseoli found on Physalis peruviana, Rumex dentatus and Solanum nigrum in Pakistan, 171
---: maize dwarf mosaic virus on corn in Iowa, 88
---: nematodes: citrus nematode (Tylenchulus semipenetrans) in Trinidad, 478; pseudo-root-knot nematode in Texas, 645; seed-gall nematode on grass in Montana, 45
---: Phytophthora cinnamomi, on loblolly pine in Louisiana, 596
---: --- fragariae race A-5 found in commercial strawberry plantings in the eastern United States, 351
---: Polyporus lucidus (Ganoderma lucidum) on honeylocust in Mississippi, 243
---: --- tomentosus spread in slash pine plantations in the southeastern United States, 527
---: powdery mildew: on mock orange in Georgia and the Southeast, 306; new records of powdery mildew fungi, 709
---: Puccinia zoysiae, in Mississippi, 770
---: Rosellinia herpotrichioides, in California, found on Douglas-fir seedlings, 249
---: root rot, of alfalfa caused by Phytophthora megasperma has been found in Canada, 451
---: rust: on zoysia grass in Alabama, 885; zoysia rust in Georgia, 839
---: sugarcane leaf scald, in Puerto Rico, first report on occurrence in one of the larger islands of the Antilles, 569
---: Uredo ignava, first report in the United States, found on bamboo in Florida, 112
---: wart disease, on subterranean clover in the United States, 797
New Hosts: alfalfa mosaic virus, first known report as causal agent for a disease (calico) of head lettuce in the field, Arizona, 629
---: apricot ring pox virus, 91 apricot varieties from foreign sources found to be hosts for, 289
---: aster yellows virus, on several grass hosts, 17
---: barley yellow dwarf virus, recovered from 5 weed species collected in Mississippi, believed to be natural summer hosts, 692
---: cucumber mosaic virus, causes mosaic disease of canna, 1st report of natural infection of canna by CMV in the United States, 899
---: Curvularia geniculata, on Sambucus chinensis, 488
---: --- verruculosa, on rice in India, 314
---: Cytospora canker, on thornless species of honeylocust, 13
---: Dalbulus maidis, on gamagrass, this is the 3rd plant species reported to be accepted by this leafhopper for development from egg to adult, 570
---: dwarfmistletoe, on Sitka spruce, found in Alaska, 921
---: Erwinia aroideae, on pear fruits and coriander leaves in India, 701
---: Fomes annosus, on Pinus patula, Abies bracteata, Artemisia tridentata, Archtostaphylos viscida and A. manzanita, found in California, 181
---: Lecanosticta acicola (conidial state of Systremma acicola), on Pinus banksiana, 208

---: Macrophomina phaseoli, on Solanum nigrum, found in Pakistan, 171
---: maize dwarf mosaic, on gamagrass, Illinois, 65
---: root-knot nematode, found on ponderosa pine in New Mexico, believed to be first report, 126; for Meloidogyne hapla, 523
---: seed-gall nematode (Anguina spp.), on western wheatgrass, 45
---: soybean cyst nematode, on seven additional weed hosts in Arkansas, 15
---: Tylenchulus semipenetrans, on Andropogon rhizomatus, 822
---: Xanthomonas alfalfae, readily infected several in greenhouse tests, 731
---: Xiphinema index, on 5 new hosts, 27
---: red thread fungus, on Bermuda grass in Mississippi in 1965-66, 525
New Jersey: 294; 351
New Mexico: 126; 175; 301
New York: 783; 877
New Zealand: 730
Nigeria: 606; 641; 828
Nilaparvata lugens: 453
Nitrogen: effect of various levels in the soil on stalk rot of corn incidence, 655
---: liquid: storage of pycniospores of Puccinia graminis secalis in, 576; storage of urediospores of Puccinia graminis tritici in, 502
North Carolina: 136; 308
North Dakota: 62
Nutrients: effects of varying combinations of N, P, K, and S on fruit rot symptoms in cranberry, 346; high and low N effects on susceptibility of chrysanthemum to Fusarium wilt, 882; nutrient factors plus tissue pH in mesocarp and epicarp of orange may be involved in preventing infection by Geotrichum candidum, 681

Oak: release of aeciospores from Cronartium fusiforme galls on slash and loblolly pine found to correlate well with appearance of susceptible young leaves of water oak in 2 spring seasons in southern Mississippi, 388
---: decline, of live, southern red, water, and post oaks in Texas associated with Cephalosporium, 75
---: Fusarium solani, isolated from oak cankers can infect young oak stems and is more virulent than strains of the fungus from tupelo, cottonwood, and yellow-popular, 160
---: fusiform rust, 432
---: heart rot, comparison of 2 methods for experimentally inoculating Nuttall oaks, 552
---: wilt: distribution extended in Arkansas, 862; evaluation of small oak bark beetle as a vector, 45; of northern pin, field studies of antibiotics and chemicals for control, 736
Oats: pathogenicity of Pythium aphanidermatum, 292
---: crown rust: adult plant resistant hybrids of CI 3034 should be searched for stem rust resistance, 631; 3000 isolates tested for pathogenicity on key differential oat varieties, 1961-1965, 935; 18 races new to North America isolated during recent surveys in Canada, 750
---: stem rust: discovery of a new and virulent race, 6AFH, in the Uniform Oat Rust Nursery, which attacks varieties with the A, B, D, E, F, and H genes for resistance, 751; new type of resistance of adult plants may be effective against all North American races, 631; physiologic races of Puccinia graminis in the United States in 1964, 332
Ohio: 951
Oidium caricae: 519
Oil: longevity and pathogenicity of Helminthosporium spp. stored under mineral oil, 321
---: spray, 1.25% and 2.85% of a light oil emulsion delayed field spread of pepper viruses but did not increase yields, 158
Oiliopsis taurica: 635
Oklahoma: 7; 599
Okra (see Hibiscus esculentus)
Onion: Botrytis allii attacking yellow Gilboa variety which is being grown for seed production in Israel, 21; bulb rotting by soilborne pathogens, especially Fusarium oxysporum, greatly reduced by alfalfa residue amendment, onion residue increased rotting, 469; disease associated with ectoparasitic nematode serious in Israel, 711
Oregon: 59
Ornamentals: 1965 survey in Georgia nurseries to determine prevalence of root rot organisms and which fungi were involved, 393
Ozone: injury to celery, first report in Connecticut and possibly in United States, 721

Pacific Islands: 892
Pakistan: 171
Papaya: powdery mildew, susceptibility of senescent leaves of fruiting plants, Hawaii, 519; relationship of virus incidence with distance and time within a field, 908
Paratylenchus curvitatus: 877
Parthenocissus tricuspidata: new host for Xiphinema index, 27
Pasmo: of flax, serious reduction of crop value in Texas, 840
Pathogenicity: of Lophophacidium hyperboreum, 356
Pea: common pea mosaic virus, strongly indicated as strain of bean yellow mosaic virus, 337
---: downy mildew, number of diseased pods reduced about 50% with 1 fungicidal spraying at the onset of flowering in western Europe, 257
---: root rot, study of chemical control using methionine and Dexon, divergence between results from field and greenhouse tests, 91
Peach (see also Prunus): 681
---: latent viruses, transmission in commercial seed, 740
---: mule's ear virus disease, 64
---: Rhizopus rot, control studies with preharvest sprays of 2,6-dichloro-4-nitroaniline (DCNA) in combination with adjuvants and with use of both hard and soft water under Colorado conditions, 565
Peanut: mycoflora isolations from peanut hypocotyls and senescent cotyledons reveals a similarity in the spectra of fungi harbored by tillering plants grown in 2 Georgia soils from fungicide-treated seed, 162
---: Aspergillus crown rot, serious outbreak in New Mexico in 1965, 175
---: bacterial wilt: disease reaction and symptom expression of 17 cultivars, 520; rapid detection of, in plant by agar-diffusion method, 836
---: leafspot, technique for forecasting, 810
---: southern blight, PCNB effective control of Sclerotium rolfsii when blight is principal disease problem in irrigated fields, 855
---: stunt, new virus disease causes severe dwarfing and fruit suppression, mechanical, graft, dodder, and aphid, transmission, 139
---: virus, severe stunting and yield reduction of 2 varieties and obvious spread of unidentified disease in 5 North Carolina locations, 136
Pear: Stewart strain of Bartlett, further tests on susceptibility to fire blight, 787
---: decline virus, new brown leaf-vein symptom associated with infected greenhouse-grown plants, 270
---: soft rot, of fruit, new host for Erwinia aroideae, India, 701
Pecan: scab, 4 new fungicide formulations tested for control in Oklahoma, 599
Penicillium digitatum: 874
--- expansum: 913
--- italicum: 874
Peniophora gigantea: 747
Pennsylvania: 6; 348; 494; 701
Peony: tree, 3 species from Japan intercepted by plant quarantine, buds were infected with the nematode Aphelenchoides fragariae, 798
Pepper: 681; seed, fungal microflora completely inhibited by new antibiotic, Pentaene G8, 299
---: viruses: new strain of tobacco mosaic virus naturally occurring on chilli pepper variety puri orange in India, virus has wide host range, 619; oil sprays delay field spread but do not increase yields, 158
Peppermint: Verticillium wilt of, most effective soil fumigants of several tested, controlled for 3 consecutive crops, 97
Peridermium stalactiforme: 144
--- harknessii: 550; 741
Peronospora pisi: 257
Pesticides: application on cranberry with "ground effect" machine shows promise, 715; common names for, 725
Phialophora malorum: 701
Phomopsis citri: 860
Phyllactinia guttata: 306
Phyllanthus acidus [=Cicca acida]: new anthracnose disease of, in India, 313
Physiological diseases: lenticel injury to apples in storage in Washington, 885
Physoderma trifolii: 796
Phytophthora spp.: 393; 695
--- cactorum: 267; 283
--- cinnamomi: 596; evidence for existence of biotypes of, based on differential pathogenic effects to the same and different broadleaf and coniferous evergreens, 647; 13 isolates from various hosts and locations differed in degree of pathogenicity to 6 Taxus cultivars, 84
--- citricola: 494
--- citrophthora: 251; 494; 867
--- fragariae: race A-5, found in two

commercial strawberry plantings for first time in eastern United States, 351
--- hibernalis: 494
--- infestans: 686
--- megasperma: 451
--- --- var. sojae: 104
--- palmivora: 422
--- parasitica: 686
--- phaseoli: 734
Picea sitchensis: first report of dwarf-mistletoe on, 921
Pinus spp.: annosus root rot, significantly higher incidence of infection revealed by core sampling than by field diagnosis, 589; cycloheximide and phytoactin evaluated for control of western gall rust and aspen cankers on lodgepole and ponderosa pines, 741; distribution and severity of Fomes annosus root rot in Illinois pine plantings, 446; fruiting of Fusarium lateritium pini on naturally infected pitch cankers of a Virginia pine in North Carolina, 476; fusiform rust, time of maximum teliospore production and weather conditions necessary for maximum infection in South Carolina, 432; incidence of Polyporus tomentosus fruiting on infected trees in a Georgia slash pine plantation, 360; loblolly pine stands given low, intermediate, or high hazard ratings for benefit of management foresters, 510; occurrence of sweet fern blister rust in Alberta and Northwest Territories could be serious threat to hard pine nurseries and plantations, 419; red pine root rot study conducted in West Virginia nursery, 545; Scleroderris lagerbergii responsible for widespread and severe dieback and mortality in red and jack pines in Upper Michigan, 402; slash and loblolly infected with Cronartium fusiforme galls serve as inoculum for infection of young water oak leaves by aeciospores in Missisissi, 388
--- banksiana: new host for brown spot needle blight caused by Lecanosticta acicola, and first report of this fungus in Canada, 208
--- contorta: comandra rust, evaluation of antibiotics for control, 528
--- --- var. latifolia: 208; white spored Peridermium stalactiforme in Alberta, Canada, 144
--- elliottii var. elliottii: development of pycnia and aecia in albino form of Cronartium fusiforme on inoculated seedlings, 842; incidence of Polyporus tomentosus in plantations in southeastern United States, 527
--- lambertiana: root disease control by soil fumigation in forest nurseries of California, 245; white pine blister rust, evaluation of basal stem application of cycloheximide and phytoactin for control in Oregon and California, 554
--- patula: 181
--- ponderosa: experimental inoculation with western gall rust for resistance studies, 550; root disease survey to determine extent of damage in natural forest stand shows that Fomes annosus is greater menace in California than past reports indicate, 439
--- --- var. scopulorum: root-knot nematode possibly associated with decline in New Mexico, believed to be first report of Meloidogyne (may be undescribed species) on this host, 126
--- radiata: Dothistroma pini on, first report of this fungus in Chile, 280
--- resinosa and P. rigida: Fomes annosus infection of stumps effectively prevented by creosote, urea, and liquid borate after 10 months, 22
--- strobus: blister rust cankers not controlled in eastern white pine by cycloheximide and phytoactin antibiotics as reported in western white pine, 224
--- taeda: Phytophthora cinnamomi and Pythium spp. associated with decline, 1st report of P. cinnamomi associated with pine in Louisiana, 596; stumps inoculated with Fomes annosus resulted in increased penetration under induced drought conditions, 747
Pistacia spp.: new hosts for Xiphinema index, 27
Plasmodiophora brassicae: 133
Pleurotus ostreatus: 552
Plum (see also Prunus): line pattern virus symptoms on indicator trees from virus-infected plum trees, 278
Podosphaera leucotricha: 645
Polyporus lucidus: 945; (Ganoderma lucidum), associated with death of honeylocust in Mississippi, 243
--- tomentosus: 360; 527
Pome Fruit Virus Subproject (IR-2): an interim report, 461
Potato: black-leg and tuber rots, investigations in Arizona revealed inter-

action between Erwinia sp. and Pythium aphanidermatum, 414; Irish, 681; seed production experiments in the United Arab Republic, 761
Powdery mildew: of apple and bean, controlled by 2 new sulfites, 645; combined control of mildew and blight of carrots, 594; on mock orange, reported for first time from Georgia and the Southeast, 306; new records of Erysiphaceae, 709; of papaya, susceptibility of senescent leaves of fruiting plants, 519; of snap bean, several fungicides tested for control, captan + tribasic copper sulfate gave excellent results, 847
Pratylenchus penetrans: 923
--- vulnus: 858
Prune (see also Prunus): bioassay of Italian variety by branch-injection indicate that 8-quinolinol phosphate and 8-quinolinol acetate warrant further investigation as systemic fungicides for orchard trees, 773
Prunus spp. (see also common names): ringspot virus complex from plum causes atypical reaction on cherry indicator tree, 278; seven viruses tested for transmission through Prunus seed, 902
Pseudomonas glycinea: 560; identification of 7 pathogenic races, 557
--- maculicola: 192
--- phaseolicola: 29; 110; 238; 371
--- solanacearum: 275; 520; 836
Pseudoperonospora cubensis: 340
Pseudopeziza ribis: 196
Pseudopityophthorus spp.: 45
Pseudotsuga menziesii: Rhabdocline needle cast, needle weight determination method accurately estimates intensity of disease, 406; Rosellinia needle blight on seedlings, first report of Rosellinia herpotrichioides in California, 249
Psylla pyricola: 270
Puccinia coronata var. avenae: pathogenicity of isolates collected in USA, 1961-1965, 935
--- graminis: method of separation and identification of isolates from mixtures on differential host varieties, 312; physiologic races in the United States in 1964, 332
--- --- f. sp. avenae: 571; 631
--- --- var. phlei-pratensis: 849
--- --- secalis: storage of pycniospores in liquid nitrogen, 576
--- --- tritici: 299; 673; storage of urediospores in liquid nitrogen, 502
--- recondita: 108
--- --- f. sp. tritici: 7th revision of International Register of Physiologic Races of Puccinia recondita f. sp. tritici, 756
--- sorghi: 119
--- striiformis: 205; 806
--- zoysiae: 839; 885; found in Mississippi, 770
Puerto Rico: 484
Purple stain: caused by Cercospora kikuchii, relation of soybean seed infection to duration of pre-flowering, flowering, post-flowering, and pod-maturity periods, 464
Pyrenochaeta terrestris: 469
Pythium spp.: 393; 596; 695; detection of in soil, 791
--- aphanidermatum: 414; demonstrated to be pathogenic on barley, oats, rye, wheat and sorghum for the first time, 292
--- debaryanum: 614
--- ultimum: 614

Quarantine: interception in 1965-66 season by plant quarantine inspectors of 3 tree peony species from Japan and of Lilium regale bulbs from the Netherlands infected by the nematode Aphelenchoides fragariae, 798; proposed isolated laboratory for study of plant pathogens, especially viruses and their vectors, 803
Quince: virus and virus-like diseases in California, 211

Races: new physiologic race of Bremia lactucae seriously affects 3 previously immune lettuce varieties in south Texas, 460; five para rubbertree clones designated as differentials for identifying races of Dothidella ulei, 187; Race 2 of Fusarium oxysporum f. lycopersici pathogenic to tomato stocks containing Acc. 160 factor for resistance, 707; serological procedures simplify differentiation of race 1 from race 2 of Fusarium oxysporum f. sp. lycopersici, 444; race 1 of Phytophthora megasperma var. sojae caused rot of soybeans on the High Plains of Texas in 1965, 104; of Pseudomonas glycinea on soybean; first report of 7 separate pathogenic races, distinguished by reactions of 7 differential varieties in Minnesota, 557; Pseudomonas solanacearum races 1, 2 and 3, serological relationships among, 275; of Puccinia sorghi, morphology and pathogenic

reactions of 2 clonal lines and their inbred recombinants on a homozygous susceptible corn, 119; of sorghums resistant to sooty stripe disease, studies indicate that frequency of resistance is associated with geographic point of origin, 606; evidence for existence of physiologic races of stripe smut attacking Kentucky bluegrass, 430; of Xanthomonas malvacearum from Australia, 323
---: rust: of flax rust in the Canadian prairies and in North Dakota in 1963 and 1964, 62; of oat crown rust, 935; 18 oat crown rust races new to North America isolated during recent surveys in Canada, 750; of oat stem, identification of a new and virulent race, 6AFH, from 4 collections of oat stem rust made in Uniform Oat Rust Nursery, 571; oat stem, in a test of select varieties and lines of oats for resistance to stem rust races, adult plants of CI 3034 was most promising source of resistance to 6AFH in hexaploid oats, 631; of Puccinia graminis, in the United States in 1964, 332; race 56 of Puccinia graminis tritici constituted more than 97% of the pathogen population in wheat in 1960-62, 90% in 1963, 673; seventh revision of the International Register of Physiologic Races of Puccinia recondita f. sp. tritici, 756; new stripe rust race pathogenic to 'Suwon 92' wheat, 205
Radiation: gamma, used on oranges in Israel to prolong storage life of fruit, 874
---: ultra-violet, did not stimulate perithecia formation in otherwise non-perithecial cultures in tests for sexual expression in Colletotrichum orbiculare, 871
Ramulispora sorghi: 606
Red thread: found on Bermuda grass, 525
Registration-certification program: organization, history, and progress of Arizona's cooperative system for development of virus-free citrus budwood, 145
Reservoir hosts: 5 Nicotiana tabacum varieties maintained highly infectious cucumber mosaic virus inoculum after 1 1/2 years, 79
Resistance: cantaloup introduction, PI 140471, approaches immunity to gummy stem blight in the field, 661
Rhabdocline pseudotsugae: 406
Rhamnus cathartica: mosaic disease on ornamental plantings found to be caused by cucumber mosaic virus, 613
Rhizoctonia spp.: 172; 393; 695
--- solani: 301; 340; 435; 484; 578; 614; 622; 852; isolated from cotton seed that were acid delinted and surface sterilized with alcoholic mercuric-chloride, 770; role in obtaining and maintaining an adequate tomato stand for production in south Florida, 124
--- leaf blight: of azalea, report on disease and control, 354
Rhizopus stolonifer: 565
Rhopalosiphum maidis: 862
--- padi: 380
Ribes alpinum: anthracnose, variation in Pseudopeziza ribis isolates from several hosts, 196
Rice: grassy stunt virus disease, transmission by the planthopper Nilaparvata lugens, 453; new host of Curvularia verruculosa, 314
Root: inhabiting fungi associated with birdsfoot trefoil, 753
--- diseases: in California forest nurseries, control by soil fumigation, 245; disease survey at Boggs Mountain State Forest shows that Fomes annosus is greater menace in California than past reports indicate, 439
--- knot: considerably reduced in Indian tomato and okra fields by addition of oil-cake amendments to the soil, 668
Rose: hybrid tea, spotted anthracnose of petals caused by Spacheloma rosarum, 934
Rosellinia herpotrichioides: 249
Rot: Aspergillus crown, of peanut, serious outbreak in irrigated fields in New Mexico in 1965, 175
---: bacterial soft rot: of Iceberg (Great Lakes) lettuce in Florida Everglades, 938; of pear fruits and coriander leaves, new hosts for Erwinia aroideae, 701
---: black core, of mandarin oranges, caused by Alternaria tenuis, 127
---: blue-mold, of apple, fungicides for control, 913
---: brown: of citrus, study of length of incubation period of Phytophthora citrophthora under natural conditions, 251; 3 Phytophthora spp. associated with brown rot syndrome in Western Australian citrus orchards, 494
---: brown stem, of soybeans in Mexico, 598
---: collar, of 1 to 7 year old apple trees propagated on East Malling II and

Malling Merton 104 rootstocks, 267
---: cotton boll, method for estimating losses, 254
---: foot, of winter wheat, fall applications of diuron to control weeds reduced incidence and severity of, 852
---: fruit, of apple, caused by Phialophora malorum, 1st report in Pennsylvania, 701
---: heart, comparison of 2 methods for experimentally inoculating Nuttall oaks, 552
---: leather, of strawberry, occurrence despite lack of rainfall attributed to heavy dew and fog conditions in central California coast, 283
---: perennial canker, of apples in West Virginia and Pennsylvania, 348
---: Phytophthora, of soybeans, on the High Plains of Texas in 1965, 104
---: Rhizopus, of peaches, preharvest spray experiments in Colorado, 565
---: root: of alfalfa in Ontario, fungi associated with, 695; Phytophthora root rot, of alfalfa, found in Canada for first time, 451; identity and relative importance of major root rot pathogens of snap beans in Wisconsin, 578; 1965 survey in Georgia to determine prevalence of organisms in ornamental nurseries and to determine which fungi were involved, 393; Fusarium, easy method for evaluating hundreds of lines and varieties of wheat at maturity for resistance, 26; of pea, study of chemical control using methionine and Dexon, divergence between results from field and greenhouse tests, 91; of pine, survey of distribution and severity in Illinois, 446; of red pine seedlings, study of nematode-fungi interaction in soil, 545; caused by Polyporus lucidus, tests of pathogenicity on important shade and timber trees of Mississippi Delta, 945; of Taxus, 13 diverse Phytophthora cinnamomi isolates varied in degree of pathogenicity to 6 cultivars, 84; of tomato, 124
---: ---: annosus, significantly higher incidence of infection of Pinus spp. revealed by core sampling than by field diagnosis, 589
---: root and butt: distribution of Fomes annosus in natural forests of California, 832; red root and butt, incidence of Polyporus tomentosus fruiting on infected slash pine trees in a Georgia plantation, 360
---: soil, of cucumber, control, 340
---: stalk: of corn, reaction of inbred lines, varieties, and hybrids to 4 causal fungi, 401; of sorghum, caused by Aspergillus niger, intensified by addition of Macrophomina phaseoli, 352
---: stem, of geranium, satisfactory technique for checking presence of bacteria in commercial plants, 515
---: stem-end, of Florida lemons for fresh fruit market, 860
---: Thielaviopsis basicola, of gloxinia rhizomes from Europe, 709
Rotenoid: concentration in Tephrosia vogelii not affected by fungicides, 484
Rotylenchulus reniformis: 149; 500
Rubber plant: "leaf stripe" disease, foliar nematode shown to be causative agent, 689
Rubbertree: (para), 5 clones designated as differentials for identifying races of Dothidella ulei, 187
Russeting: of McIntosh and Cortland apples, possibly physiological disease, 151
Rust: first report of Uredo ignava in the United States, found on bamboo in Florida, 112; morphological evaluation of 2 pathogenic clonal lines of Puccinia sorghi and their inbred recombinants on a homozygous susceptible corn, 119; seventh revision of the International Register of Physiologic Races of Puccinia recondita f. sp. tritici, 756
---: blister, of eastern white pine, not controlled in the Lake States by cycloheximide and phytoactin, contary to results on western white pine, 224
---: comandra, on lodgepole pine, evaluation of antibiotics for control, 528
---: crown, of oat, pathogenicity of isolates collected in USA, 1961-1965, 935; 18 oat crown rust races new to North America isolated during recent surveys in Canada, 750
---: flax, 16 races in varying frequencies in Canadian prairies and in North Dakota, 1963 and 1964, 62; serious reduction of crop value in Texas, 840
---: fusiform: development of pycnia and aecia in albino form on pine, 842; sporophores of Polyporus tomentosus var. circinatus associated with basal cankers caused by fusiform rust fungus, 527; study of conditions essential to teliospore germination and sporidial production in South Carolina, 432

---: leaf: effect of inhibiting growth of 14 wheat varieties with Cycocel on leaf rust infection types, 108; of timothy, 849
---: --- and stem, of timothy, in field and greenhouse screening of 100 Phleum introductions, selections out of PI's 251846, Italy and 267063, Poland were completely resistant -- good resistance found in selections out of 16 other accessions, 849
---: stem: of oats, a new and virulent race, 6AFH, important because virulence for all described genes has been combined into a single phenotype, 571; of oats, varietal reaction to races 6AF and 6AFH reveals C I 3034 as new source of resistance to 6AFH in adult plants in Minnesota, 631; physiologic races of Puccinia graminis in the United States in 1964, 332; of timothy, 849; of wheat, inhibition by new antifungal antibiotic, 299
---: stripe: of grasses, systemic activity of 2 new fungicides, 624; new race pathogenic to 'Suwon 92' wheat, 205; of wheat, effective chemical seed treatment for systemic control of seedling infection, 806
---: sweetfern blister: amendments to records of distribution and hosts, 744; occurrence in Alberta and Northwest Territories could be serious threat to hard pine nurseries and plantations, 419
---: western gall, antibiotic tests for control on lodgepole and ponderosa pines, 741; experimental inoculation of ponderosa pine for resistance studies, 550
---: wheat, rapid method of assaying pathogenic potential of populations of Puccinia graminis tritici, 673
---: wheat stem, evaluation of nickel + dithiocarbamates for control, 373
---: wheat stripe, mist culture apparatus used for root development studies, 369
---: white pine blister, of sugar pine in Oregon and California, evaluation of basal stem application of cycloheximide and phytoactin for control, 554
---: white spored Peridermium stalactiforme, on lodgepole pine in Alberta, Canada, 144
---: zoysia: found in Alabama for first time, 885; in Georgia, 839
Rye: pathogenicity of Pythium aphanidermatum, 292
Sambucus chinensis: new host for Curvularia geniculata, 488
Sanitation: 10 recommendations for control of angular leaf spot of cotton that successfully eradicated Xanthomonas malvacearum from California, 168
Scab: pecan, four new fungicide formulations tested for control, 599
Scleroderris lagerbergii: 402
Sclerospora sorghi: 641
Sclerotinia borealis: 69
--- camelliae: 525
--- homeocarpa: 770
Sclerotium bataticola: 401
--- rolfsii: 443; 855
Seed: potato, production experiments in U.A.R., 761; Rhizoctonia solani isolated from cotton seed, 770; size of Funza barley seed has significant effect on smut percentage, 101
---: transmission: of latent viruses in commercial peach seed, 740; of psorosis virus through citrus hybrid at levels of 15 to 31%, 81; western X-disease of chokecherry not definitely found to be carried in seed, however nursery seed should come from healthy trees since only low percentage from infected trees will germinate, 659
--- treatments: effective chemical treatment for systemic control of seedling infection of wheat by stripe rust, 806; effect on soil-borne fungi at the hypocotyl and cotyledonary surfaces of peanut seed, 162; fungicide evaluations for control of damping-off of table beet seedlings indicate that if PCNB is combined with a standard material used for control of Pythium diseases, Rhizoctonia in the soil, which varies greatly from year to year will also be controlled, 614
Septoria linicola: 840
Serodiagnosis: use of disposable leaf presses for obtaining plant sap for, 822
Serological studies: relationships among races 1, 2 and 3 of Pseudomonas solanacearum and 4 other plant pathogenic bacterial species, 275
Serum: anti-, against Corynebacterium michiganense can aid in detection of tomato bacterial canker, 96; anti-, against Pseudomonas solanacearum races 1, 2 and 3 and a study of their relationships, 275; anti-, against several isolates of Pseudomonas solanacearum and the serological

detection of the fungus in peanut plants by a rapid agar-diffusion method which distinguishes bacterial wilt from wilts caused by other organisms, 836
Sexual expression: in Colletotrichum orbiculare, 871
Slugs: transmission of cabbage leaf spot caused by Alternaria brassicicola by, in France, 764
Smut: increase in susceptibility of corn inbred W22 by adding potassium to soil associated with decrease in reducing sugar and increase in nitrogen in the pith tissue, 12; survival of various smut fungi cultures for 11 years under mineral oil, 800
---: flag and stripe, systemic activity of 2 new fungicides, 624
---: head, of corn in Idaho, 215
---: stripe: of Kentucky bluegrass, evidence for existence of physiologic races, 430; prevalence, pathogenicity and control studies in New Jersey turf and forage grasses, 294
Snow mold: on 4 bluegrass varieties, 3 chemicals tested for control during 3 winter seasons in Alaska, 69; further evidence that Fusarium nivale is a primary root-rot pathogen of cereals and that in sequence with Bipolaris sorokiniana it causes damage in warm as well as in cold seasons in winter wheat in Washington, 154
Soil: detection of Pythium in, 791; induced moisture soil effect on growth of Fomes annosus in inoculated loblolly pine stumps, 747; isolating Verticillium from, 897; textural studies of in relation to development of hazard ratings for Fomes annosus in pine plantations in Virginia, 510
---: amendments, for the control of onion infection and bulb rotting, 469
---: fumigants, for control of Verticillium wilt of peppermint, most effective treatments controlled for 3 consecutive crops, 97
---: treatments: combinations tested for control of 2 Heterodera spp. occurring with club root organism in Brussels sprouts, 133; effects of chemicals and mulching on development of new apple trees when used at planting time, 201; root disease control by fumigation in California forest nurseries, 245
Sooty stripe disease: testing for resistant races of sorghums in Nigeria, 606
Soreshin: 124
Sorghum: Colletotrichum graminicolum spreads through dead stalk tissue cells and is inhibited by living cells, 287
---: downy mildew, part of world collection and Nigerian collection of grain sorghums screened for resistance in 1964 and 1965, 641
---: ergot, host range and epidemiology in Nigeria, 828
---: Pythium aphanidermatum, pathogenicity of, 292
---: sooty stripe, studies in Nigeria of the world collection of sorghum indicate that frequency of resistance is associated with geographic point of origin, 606
---: stalk rot, caused by Aspergillus niger, intensified by addition of Macrophomina phaseoli, 352
South Carolina: 432; 482
South Dakota: 819; 917
Soybean: bacterial blight, inoculation procedures for comparing reaction of soybeans to pathogen, 560
---: brown stem rot, in Mexico, 598
---: cyst nematode, comparative rates of reproduction on 12 hosts, 608
---: Phytophthora rot, found on the High Plains of Texas in 1965, 104
---: Pseudomonas glycinea, 7 pathogenic races identified in Minnesota, 557
---: purple stain in seed coats found to be closely related to length of flowering stage in selected commercial varieties and plant introductions, 464
---: tobacco ringspot virus of, losses caused by, 910
Sphacelia sorghi: 828
Sphaceloma rosarum: 934
Sphacelotheca spp.: 800
--- reiliana: 215
Spore trap: simple, inexpensive, continuous sampling, 906
Squash (see Cucurbits)
Stalk rot: of corn, there was little difference between 3 rating methods evaluated and disease means were not much affected by high N fertilization levels, 655
Stem canker: oak strain of Fusarium solani can infect young oak stems and is more virulent than other strains tested, 160
Stemphylium botryosum f. lycopersici: 635
Stinkhorn fungus: unusual occurrence on golf turf, 840
Storage: lenticel injury to apples resulting

from fruit packed in molded pulp trays containing urea-formaldehyde wet-strength resins, 885; of pycniospores of Puccinia graminis secalis in liquid nitrogen, 576; storage period of oranges extended in Israel by gamma radiation which either delayed incubation of Penicillium fungi responsible for decay or prevented their development, 874; of urediospores of Puccinia graminis tritici in liquid nitrogen, 502
Strains: evidence that common pea mosaic virus is a strain of bean yellow mosaic virus, 337; of maize dwarf mosaic virus, 2 distinct strains in northeastern United States identified by host reaction and serological relationships, 814; new strain of tobacco mosaic virus in India, 612; of Rhizoctonia solani and Thielaviopsis basicola which cause cotton seedling diseases, relation between pathogenicity and sensitivity to chemicals, 301
Strawberry: new disorder of fruit, caused serious losses in Salinas-Watsonville district of California, 327
---: angular leaf spot, found in Wisconsin for first time, 167
---: leaf spot, relative resistance of 63 varieties and selections studied at 2 locations for 2 years, 105
---: leather rot, occurrence during the absence of rainfall attributed to heavy dew and fog conditions combined with cool temperatures in central California coast, 283
---: Phytophthora fragariae, race A-5 found in Maryland and New Jersey, this is the first report in commercial strawberry plantings in the eastern United States, 351
---: viruses: rapid spread into new plantings in western Washington, 586; crinkle virus, over 50 chemical compounds tested did not inactivate virus when sprayed on foliage of infected alpine hybrid vesca plants, 843; virus infection indexing, a simple, efficient, and inexpensive method, 426
Streptomyces anandii: 299
Sugarbeet: nematodes, evaluating varieties and breeding lines for tolerance to nematode invasion, no outstanding resistance was observed, 650
---: viruses: cucumber mosaic virus infectivity increased in vitro on infected leaf tissue, 639; savoy and yellow vein viruses in Great Plains, 1963-64-65, root weight and percentage of sucrose significantly reduced, 457
Sugarcane: leaf scald, found in Puerto Rico, first report of occurrence in one of the larger islands of the Antilles, 569
Sunflower: Mexican, wilt of, caused by celery-wilt Fusarium, 391
Surfactants: effect on fungicides applied to watermelon foliage, 327
Sweetgum: 945
Sweetpotato: internal root necrosis of New Zealand's 'Owairaka Red', 730; varietal reaction of 24 selections to reniform nematode, 500
Systremma acicola: 208

Tanzania: 593
Tar spot: of citrus, second report in Florida, chemical control, 357
Taxus: 13 diverse Phytophthora cinnamomi isolates varied in degree of pathogenicity to 6 cultivars, 84
Teliospore: germination of Cronartium fusiforme, 432
Temperature: of citrus groves, influence on efficacy of hot-water treatment for Phytophthora citrophthora control of fruit rot, 867; effect of low postinoculation temperature on colonization of lima bean stems by Phytophthora phaseoli, 734; a factor in lenticel injury in apples, 885; hot-water treatments effective in eliminating Pratylenchus vulnus from grapevine roots, however soil treatments should still be used, 858; Thielaviopsis basicola development enhanced by temperature stress on roots of greenhouse-grown birdsfoot trefoil, 825
Tennessee: 427
Tephrosia vogelii: field evaluations at Mayaguez, Puerto Rico of 8 fungicides for effect on stand count, plant height, dry weight of leaves and stems, leaf:stem ratio, and rotenoid concentration in leaves, 484
Texas: 7; 75; 104; 149; 460; 645; 840
Thielaviopsis basicola: 301; 709; growth on birdsfoot trefoil in the greenhouse increased by temperature stress, 825
Timothy: screening of introductions for sources of resistance to stem and leaf rust among Phleum species, 849
Tobacco: diseases in Kentucky in 1965, 218
---: cyst nematodes, control with volatile and nonvolatile nematocides, 507
---: false broomrape, field observations

on response of Maryland tobacco to, 627
---: viruses: cucumber mosaic virus, highly infectious inoculum obtained from Kentucky 57, 304-1, T.I. 245, Holmes Samsoun and Xanthi-nc varieties after 1 1/2 years, 79; survey of virus diseases for distribution and relation of previous cropping history to occurrence in North Carolina, 308
Tomato: 681; unusual occurrences of late blight, buckeye rot, leaf mold and bacterial spot in Louisiana, 686
---: bacterial canker, 3 serological diagnostic methods for, 96
---: bacterial spot, analysis of role in production in south Florida has resulted in effective control program using copper, maneb, and streptomycin, 699
---: curly top virus, wooly and dwarf tolerance to, Colorado, 309
---: foliage diseases, severity affected by mode of irrigation in Israel, 635
---: Fusarium oxysporum f. sp. lycopersici, serological procedures simplify differentiation of race 1 from race 2, 444
---: nematodes, 'Rutgers', used as indicator in nematocide dose and exposure time tests, 915
---: Rhizoctonia solani, role in obtaining and maintaining an adequate tomato stand for production in south Florida, 124
---: southern blight, losses reduced, and fruit yields and plant vigor improved by deep plowing, 443
---: wilt, every major tomato-growing area of Florida resurveyed for race 2 of Fusarium oxysporum f. lycopersici which is pathogenic to stocks containing Acc. 160 factor for resistance, 707
Top damping-off: of tomato, 124
Toxin: production in vitro by non wilt-inducing isolates of Fusarium, 261
Transmission: apple virus transmission studies with selective Malus indicators, 40; bud transmission of dwarfing virus in sweet cherry, 367; of cabbage leaf spot by slugs in France, 764; graft, of a ringspot-like virus from white ash to herbaceous plants, 905; of latent viruses in commercial peach seed, 740; of seven viruses through Prunus seed tested, 902
Trichoderma viride: 747
Trinidad: 478
Tung: angular leaf spot: effect on yield and oil content of fruit from trees sprayed with Bordeaux mixture or petroleum oil for 5 years, 54; evaluation of various fungicides for control, 717
Tylenchulus semipenetrans: 822; 960

United Arab Republic: 761
Uredo ignava: 112
Ustilago spp.: 800
--- maydis: 12
--- nuda: 101
--- striiformis: 294; 430

Vectors associated with: barley yellow dwarf virus, on wheat -- Rhopalosiphum padi, 234
---: corn stunt virus, -- corn leafhopper, gamagrass found to be new host for, 570
---: cucumber mosaic virus, of Rhamnus cathartica -- Myzus persicae, 613
---: decline virus of pear -- pear psylla, 270
---: grassy stunt disease of rice,-- planthopper, Nilaparvata lugens, 453
---: maize dwarf mosaic virus of corn -- corn leaf aphid, 862
---: oak wilt -- small oak bark beetle, 45
---: strawberry viruses -- strawberry aphid, 586
---: streak mosaic, of wheat -- Aceria tulipae, 917
---: stunt (new virus disease) of peanut -- green peach aphid, 139
---: watermelon mosaic virus -- melon aphid, 305
---: wheat streak mosaic virus: of corn -- Aceria tulipae, 819; of corn and wheat -- wheat curl mites, 951
Verticillium spp.: method for isolating Verticillium from field soil, 897
--- albo-atrum: 930
--- dahliae (=V. albo-atrum): 97
Vigna sesquipedalis: intercellular colonization and the production of pectic enzymes by Malayan isolates of Fusarium, 116
Vinca minor: dagger nematodes can cause severe decline under some circumstances, 420
Virginia: 139; 510
Virus, Virus disease: availability of "Index of Plant Virus Diseases", 624; establishment of virus-free citrus budwood through Arizona's Cooperative Citrus Registration-Certification Program, 145; possible cause of an internal root necrosis of a New Zealand sweetpotato, 730;

first report of a virus resembling tobacco streak in western United States isolated from dodder growing on a desert shrub in southern California, shrub may be symptomless reservoir for virus, 479
---: of apple: preliminary report of IR-2 Pome Fruit Virus Subproject stating objectives and procedures necessary for virus indexing of latent diseases, 461; transmission studies with selective Malus indicators, 40
---: apricot ring pox, 91 foreign varieties found to be hosts, 289
---: aster yellows, susceptibility of 44 grasses tested, host range now includes 8 grasses (including oats), 17
---: barley yellow dwarf: report of recovery tests on 10 weed species from Mississippi suspected to be natural summer hosts for BYDV, 692; resistance in selected barley varieties, 380; 12 wheat varieties susceptible to, in Kansas, 234
---: of beans: tests of bean cultivars and breeding lines for sources of resistance to bean yellow mosaic, alfalfa mosaic, western ringspot, and common bean mosaic viruses, 532; common pea mosaic strongly indicated as strain of bean yellow mosaic, 337
---: calico, of lettuce, caused by alfalfa mosaic not by tobacco ringspot, 629
---: of Cattleya, simple method for detection of a flexuous-rod virus associated with infectious Cattleya blossom necrosis, 779
---: of cherry, bud transmission of dwarfing virus in sweet cherry, 367
---: common pea mosaic, strongly indicated as strain of bean yellow mosaic, 337
---: of corn: distribution and identification of mechanically transmissible corn viruses in Louisiana, 325; occurrence of mechanically transmissible virus of corn in Mississippi, 409; evidence that streak disease of corn in California is caused by sugarcane mosaic and that corn stunt does not occur in that State, 318; corn stunt, corn leafhopper found on new host, gamagrass, in Mississippi, 570
---: of cranberry, virus-like ringspot symptoms on fruits and leaves of Searles and Howes varieties in Wisconsin, 543
---: crinkle, of strawberry alpine hybrid vesca, studies on inactivation by chemotherapy, 843
---: cucumber mosaic: causes mosaic disease of canna, 899; causes mosaic disease of Rhamnus cathartica in Wisconsin, 613; 5 Nicotiana tabacum varieties serving as reservoir hosts maintained highly infectious inoculum after 1 1/2 years, 79; from sugarbeet, in vitro increase in virus infectivity, 639
---: cucurbit viruses, on the west coast of Mexico, 947
---: curly top, tolerance of woolly and dwarf tomatoes to, 309
---: grassy stunt, of rice, transmission by planthoppers, 453
---: green ring mottle virus, of sour cherry, severe fruit necrosis caused by strains of, 59
---: maize dwarf mosaic: a heavy-duty, mobile air blast inoculator has been designed, built, tested, and proved to be rapid and efficient for both field and greenhouse use, 363; 2 distinct strains occurring in northeastern United States, 814; presence established in several counties in Pennsylvania in 1965, 6
---: ---: of grasses: mechanical inoculation of St. Augustine grass seedlings resulted in 80% infection, 441; first report of maize dwarf mosaic on gamagrass, Illinois, 65; positive transmission of one isolate from Johnson grass in Kansas, first report, 1965, 11
---: ---: of corn: found in Iowa for first time, 88; host range, properties, and partial purification of a Missouri isolate, 862; in Tennessee, 427
---: of Malvaceous plants, chemo- and thermotherapy for control of leaf curl, 88
---: mule's ear, of peach, 64
---: muskmelon necrotic fleck, new strain of tobacco mosaic virus isolated from muskmelon in Arizona, 49
---: of papaya, relationship of disease with distance and time within a field, 908
---: of peach, transmission of latent viruses in commercial peach seed, 740
---: of peanuts: new disease (stunt) causes severe dwarfing and fruit suppression, mechanical, graft, dodder, and aphid transmission, 139; severe stunting and yield reduction, and obvious spread of unidentified virus in 5 North Carolina locations, 136

---: pear decline, new brown leaf-vein symptom associated with infected greenhouse-grown pear plants, 270
---: of pepper, oil sprays delay field spread but do not increase yields, 158
---: phloem necrosis, (Morsus ulmi) of American elm in Mississippi Delta, 751
---: potato, seed production experiments in U.A.R., 761
---: of Prunus: 7 viruses tested for transmission through Prunus seed, 902; Prunus ringspot complex from plum causes atypical symptoms on Shiro-fugen flowering cherry, 278
---: psorosis virus, transmitted through seeds of at least one citrus hybrid ('Carrizo' citrange) at levels of 15 to 31%, 81
---: of quince, in California, 211
---: sour cherry yellows, of cherry, effect of gibberellic acid sprays on fruit spur development of infected trees at different growth stages and environmental conditions, 240
---: of strawberry: rapid spread into new plantings in western Washington, 586; a simple, efficient, and inexpensive method of collecting leaves for virus infection indexing, 426
---: of sugarbeet, savoy and yellow vein diseases significantly reduced root weight and sucrose percentage in Great Plains, 1963-65, 457
---: sugarcane mosaic, reaction of 180 sweet and dent corn varieties to Johnson grass strain of, 793
---: of tobacco, viruses present, relation of occurrence to previous cropping history and distribution in North Carolina, 308
---: tobacco mosaic, new strain naturally attacks chilli pepper variety Puri orange in India, and has a wide host range, 619
---: tobacco ring spot: of guar, new occurrences in Texas and Oklahoma, 7; of soybean, losses caused by, 910
---: watermelon mosaic, aluminum foil for repellency to melon aphids, 305
---: western X-disease, chokecherry infected in central and northern Great Plains, study of possible seed transmission of the disease, 659
---: wheat streak mosaic: epidemiology in South Dakota, 1962-66, and spread of virus under field and greenhouse conditions, 917; tests indicate that Ohio 3A virus culture is strain of wheat streak mosaic and its occurrence in Ohio suggests that wheat and susceptible corn could be year-round hosts throughout corn belt, 951; of corn, host range and spread of virus and vector in South Dakota, 1962-66, 819
---: of white ash, graft transmission of a ringspot-like virus from white ash to herbaceous plants, 905

Wart disease, of subterranean clover caused by Physoderma trifolii, first report on this host in the United States, 797
Washington: 154; 586; 885
Watermelon (see Cucurbits)
Weather: freeze of November 1955 in Pacific Northwest may be predisposing factor in severe dieback of Ceanothus spp., 399; frost injury on wheat, 928; temperature-relative humidity-time technique for forecasting Cercospora leafspot of peanut, 810
West Virginia: 348; 545
Wheat: frost injury, 928; pathogenicity of Pythium aphanidermatum, 292
---: Cephalosporium stripe disease, first published report from Michigan, 190
---: rots: common root rot, easy method for evaluating hundreds of lines and varieties at maturity for resistance, 26; fall applications of diuron to control weeds reduced incidence and severity of foot rot of winter wheat caused by Cercosporella herpotrichoides and Rhizoctonia solani, 852; further evidence that Fusarium nivale is a primary root-rot pathogen of cereals and that in sequence with Bipolaris sorokiniana it causes damage in warm as well as in cold seasons in winter wheat in Washington, 154
---: rusts: rapid method of assaying pathogenic potential of Puccinia graminis tritici populations for use in epidemic prediction and in detection of new, dangerous virulences, 673; seventh revision of the International Register of Physiologic Races of Puccinia recondita f. sp. tritici, 756
---: ---: leaf, effect of host growth inhibition by Cycocel on infection types, 108
---: ---: stem: evaluation of nickel + dithiocarbamates for control, 373; physiologic races of Puccinia graminis in the United States in 1964,

332; pustules markedly reduced by new antibiotic, Pentaene G8, 299
---: ---: stripe: effective chemical seed treatment for systemic control of seedling infection, 806; mist culture apparatus used for root development studies, 369; new stripe rust race pathogenic to 'Suwon 92', 205
---: viruses: barley yellow dwarf virus, field tests indicated susceptibility of 12 varieties of winter wheat in Kansas, 234; wheat streak mosaic virus in the Ohio Valley, 951; streak mosaic, epidemiology of virus in South Dakota, 1962-66, and spread of virus under field and greenhouse conditions, 917
Wilt: of Mexican sunflower, caused by celery-wilt Fusarium, 391; of alfalfa in Ontario, fungi associated with, 695
---: bacterial: of peanut, disease reaction and symptom expression of 17 cultivars, 520; Pseudomonas solanacearum, detected serologically in peanut plants by an agar-diffusion method, 836; races 1, 2, and 3 detected serologically and compared, 275
---: Fusarium: of chrysanthemum, Yellow Delaware cultivar under high N 3 weeks before inoculation more susceptible to wilt than plants under low N, 882; resurvey of every major tomato-growing area of Florida for, 707
---: oak: distribution extended in Arkansas, 862; evaluation of small oak bark beetle as a vector, 45; field evaluation of antibiotics and chemicals for control in northern pin oaks, 736
---: Verticillium: comparison of virulence of isolates of V. albo-atrum from several of the cotton-growing regions of the U.S., Mexico, and Peru, 930; of peppermint, control by soil fumigation for 3 consecutive crops, 97
Wisconsin: 167; 192; 539; 543; 578; 677; 957

Xanthomonas albilineans: 569
--- alfalfae: 731
--- fragariae: 167
--- malvacearum: 323; eradicated from California through sanitation practices, 168
--- pelargonii: 515
--- phaseoli: 29; 110; 371
--- pruni: 112
--- vesicatoria: 635; 686; 699
Xiphinema americanum: 420
--- index: 27

CROPS RESEARCH DIVISION

Prepared by Sheila D. Thompson

For sale by the Superintendent of Documents, U. S. Government Printing Office, Washington, D. C. 20402 - Price $4.00 per year, domestic; $5.00 per year, foreign. Single copy price will vary.

UNITED STATES DEPARTMENT OF AGRICULTURE
Agricultural Research Service
Beltsville, Maryland 20705

Official Business

Postage and Fees Paid
U.S. Department of Agriculture